



eduresourceza.com

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

**MARK SCHEME for the May/June 2011 question paper
for the guidance of teachers**

0580 MATHEMATICS

0580/21

Paper 2 (Extended), maximum raw mark 70

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2011	0580	21

Abbreviations

cao	correct answer only
cso	correct solution only
dep	dependent
ft	follow through after error
isw	ignore subsequent working
oe	or equivalent
SC	Special Case
www	without wrong working

Qu.	Answers	Mark	Part Mark
1	847	1	
2	correct regions shaded	1, 1	
3	48	2	B1 for 3 and 16 seen
4	(a) 10	1	
	(b) 5.5 oe	1	
5	(a) 86400	1	
	(b) 8.64×10^4	1ft	
6	108	2	M1 for 3^3 or 27 or $\left(\frac{1}{3}\right)^3$ or $\frac{1}{27}$ seen
7	13	3	B1 for 12, 5 seen M1 for $(\text{their } 12)^2 + (\text{their } 5)^2$ or M2 $\sqrt{[(-8-4)^2 + (1-6)^2]}$ oe or M1 if $\sqrt{\quad}$ missing
8	6.70	3	M1 for $(r^3 =) 1260 \times \frac{3}{4\pi}$ oe seen M1 for $\sqrt[3]{\quad}$ of their r^3 seen or implied
9	22.5 oe	3	B2 $180 = 5x + 2x + x$ oe or better B1 for $2x$ or $6x$ marked in the correct place on the diagram.
10	$x = 13$ $y = -9$	3	M1 for consistent multiplication and addition/subtraction A1 for $x = 13$ or A1 for $y = -9$
11	(a) 85.8	2	M1 for 23.25 and 19.65 seen
	(b) 456.8625 cao	1	
12	(a) (0)8(.)01 (am)	1	Not 8.01pm
	(b) 78.4 or 78.38 to 78.39	3	M2 for $827 \div 10.55$ or M1 for figs $827 \div$ their time
13	(a) 0.54	2	M1 for $\frac{2.7 \times 20000}{100000}$ oe or SC1 for figs 54 in answer
	(b) 1.61	2	SC1 for figs 161 or M1 200^2 or $20\,000^2$ seen

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2011	0580	21

14	-2.64, 1.14 cao with working	4	B1 for $\sqrt{3^2 - 4(2)(-6)}$ or better seen anywhere B1 for $p = -3$ and $r = 2 \times 2$ or better as long as in the form $\frac{p + \sqrt{q}}{r}$ or $\frac{p - \sqrt{q}}{r}$ After B0B0, SC1 for -2.6 or -2.637(45...) and 1.1 or 1.137(45...)
15	(a) 4 (b) (i) $\frac{12}{36}$ oe 0.333 (ii) $\frac{11}{36}$, 0.306 or 0.3055 to 0.3056 (c) $\frac{8}{15}$ oe 0.533(3...)	1 1 1 1	
16	(a) Answer given (b) $k = (\pm)\sqrt{\frac{4A}{(4-\pi)}}$ or $2\sqrt{\frac{A}{(4-\pi)}}$	2 3	M1 $(A =)k^2 - \pi\left(\frac{k}{2}\right)^2$ E1 $A = k^2 - \frac{\pi k^2}{4}$ correctly completed to $4A = 4k^2 - \pi k^2$ M1 factorising (must contain a π) M1 division (by coefficient of k^2) M1 square root
17	(a) 66° (b) 33° (c) 123°	2 1 2	M1 for 90° clearly identified as A B1 for OBA or $OAB = 57^\circ$
18	(a) (i) $-\mathbf{r} + \mathbf{q}$ or $\mathbf{q} - \mathbf{r}$ (ii) $\frac{1}{2}(3\mathbf{q} - \mathbf{r})$ oe (b) correct working	1 1 3	Must be simplified M1 for $\mathbf{MX} = \frac{1}{2}\mathbf{r} + \frac{3}{4}$ their $(-\mathbf{r} + \mathbf{q})$ M1 using a different route for $\mathbf{XS}$ or $\frac{1}{2}\mathbf{MS}$ E1 dep correct simplification and conclusion
19	(a) 480 (b) 9900 (c) 0.125 or $\frac{1}{8}$	1 3 2	M1 for attempt at area under graph M1 for $0.5 \times 15 \times (\text{their } \mathbf{(a)}) + 14 \times 60$ oe or $0.5 \times 15 \times (8 + 14)$ oe M1 for numerical vertical/horizontal or numerical use of $v = u + at$ but $t \leq 120$ or $t \leq 2$
20	(a) (i) 9 (ii) $8x^3$ cao (b) 4 www (c) $\frac{x+3}{2}$	1 1 3 2	M1 for $(2x - 3)^3 = 125$ M1 $2x - 3 = 5$ M1 for $x \pm 3 = 2y$ or $x = \frac{y \pm 3}{2}$