



eduresourceza.com



Education

KwaZulu-Natal Department of Education
REPUBLIC OF SOUTH AFRICA

MATHEMATICS P2

PREPARATORY EXAMINATION

SEPTEMBER 2016

MEMORANDUM


**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MARKS : 150

This memorandum consists of 15 pages.

SECTION A**QUESTION 1**

1.1 min = 9 ; maximum = 92 ; upper quartile = 75 Lower quartile = 28 and medium = 55 Therefore five number summary is (9; 28; 55; 75; 92)	A✓9 and 92 A✓28 A✓55 A✓75 (4)
1.2 	CA✓ correct Q1 & Q3 CA✓ median correctly shown CA✓ both correct whiskers (3)
1.3 Data is skewed to the left /Data is negatively skewed	CA CA✓✓ Answer (2)
1.4 mean mark for M-cee-nai High $= \frac{9+14+14+19+21+23+33+35+37+37+42+45+55+56+57+59+68+75+75+75+77+78+80+81+92}{25}$ $\frac{1257}{25} = 50,28$ OR $\bar{x} = \frac{1257}{25} = 50,28$	A✓sum CA✓answer (2) [Answer only full marks]
1.5 Bee Vee High School. Bee Vee High School performed better because half of the learners got above 60% whilst half of M-cee-nai learners got more than 55%. The median of Bee Vee High was higher than that of M-cee-nai High.	CA✓Bee Vee High CA✓✓Reasoning (3)
	[13]

QUESTION 2

2.1	(6; 160)	A✓ 6 A✓ 160 (2)
2.2	$y = -1,64x + 73,52$	AA✓✓ gradient AA✓✓ y - intercept (4)
2.3	$r = -0,2$	AA✓✓ answer (2)
		[8]

QUESTION 3

3.1	$AC = \sqrt{(-5-7)^2 + (1-(2))^2}$ $= \sqrt{(12)^2 + (3)^2}$ $= \sqrt{144 + 9}$ $= \sqrt{153}$ $= 12,37$	A✓ correct Subst CA ✓ answer (2)
3.2	$M_{BC} = \frac{6-(2)}{1-7}$ $= \frac{8}{-6}$ $= \frac{-4}{3}$ $y - y_1 = m(x - x_1)$ $y - 6 = -\frac{4}{3}(x - 1)$ $3y - 18 = -4x + 4$ $3y = -4x + 22$	A✓ $-\frac{4}{3}$ CA✓ correct subst. of (1;6) And (7;-2) CA✓ equation in any form (3)

3.3 $\hat{B} = \theta = \alpha - \beta$Ext $\angle$ $\tan \alpha = m_{BC} = -\frac{4}{3}$ $\therefore \alpha = 126,9^\circ$ $\tan \beta = m_{AB} = \frac{5}{6}$ $\therefore \beta = 39,8^\circ$ $\theta = \alpha - \beta$ $\theta = \alpha - \beta$ $= 126,9^\circ - 39,8^\circ$ $= 87,1^\circ$ $\therefore \hat{ABC} = 87,1^\circ$ OR $\text{Distance AB} = \sqrt{(1+5)^2 + (6-1)^2}$ $= \sqrt{61}$ $\text{Distance BC} = \sqrt{(1-7)^2 + (6+2)^2}$ $= \sqrt{100}$ $= 10$ $\text{Distance AC} = \sqrt{(-5-7)^2 + (1+2)^2}$ $= \sqrt{153}$ $\cos \hat{B} = \frac{a^2 + c^2 - b^2}{2ac}$ $= \frac{10^2 + (\sqrt{61})^2 - (\sqrt{153})^2}{2(10)(\sqrt{61})}$ $= 0,051$ $\hat{B} = 87,1^\circ$	CA ✓ $\tan \alpha = -\frac{4}{3}$ CA ✓ $\alpha = 126,9^\circ$ A ✓ $\tan \beta = \frac{5}{6}$ CA ✓ $\beta = 39,8^\circ$ CA ✓ $\hat{ABC} = 87,1^\circ$ (5) A ✓ Distance AB A ✓ Distance BC A ✓ Distance AC CA ✓ substitution in cosine rule CA ✓ answer (5)
3.4 $P\left(\frac{-5+1}{2}; \frac{1+6}{2}\right)$ $P\left(-2; \frac{7}{2}\right)$	AA ✓ ✓ both co-ordinates (2)



3.5 $m_{AC} = \frac{-2-1}{7+5}$ $= \frac{-3}{12}$ $= \frac{-1}{4}$ through $(-1; 3)$ equation: $y - 3 = -\frac{1}{4}(x + 1)$ $y - 3 = -\frac{1}{4}x - \frac{1}{4}$ $\therefore y = \frac{-1}{4}x + 2\frac{3}{4} \text{ or } y = -\frac{1}{4}x + \frac{11}{4} \text{ or}$ $4y + x - 11 = 0$	A ✓ $\frac{-1}{4}$ CA ✓ subst. $(-1;3)$ CA ✓ equation.in any form (3)
3.6 $m_{AB} = \frac{5}{6}; 6x + 5y = 18$ $5y = -6x + 18$ $y = \frac{-6}{5}x + \frac{18}{5}$ $\therefore m_1 = \frac{-6}{5}$ $m_{AB} m_1 = -1$ $\therefore m_{AB} \perp 6x + 5y = 18$	A ✓ $m_1 = -\frac{6}{5}$ A ✓ $m_{AB} \cdot m_1$ A ✓ $= -1$ (3)
	[18]

QUESTION 4 4.1.1 At W, $y = 2$ $3x + 4(2) + 7 = 0$ $3x = -15$ $x = -5$ W(-5;2) $r = 5$ $(x + 5)^2 + (y - 2)^2 = 25$	A✓ subst $y = 2$ CA✓ $x = -5$ CA✓ co -ordinates of W CA✓ $r = 5$ CA✓ equation of the circle. (5)
4.1.2 $VZ = 2r = 2 \times 5 = 10$ units	CA✓ answer (1)
4.1.3 $m_{GZ} = \frac{2+1}{0+1}$ $= 3$	A✓ substitution into formula CA✓ answer (2)
4.1.4 Midpoint of GZ is $\left(-\frac{1}{2}; \frac{1}{2}\right)$	A✓ coordinates (1)
4.1.5 $m_{GZ} = 3$ $m_{\perp} = -\frac{1}{3}$ $y - \frac{1}{2} = -\frac{1}{3}\left(x + \frac{1}{2}\right)$ $y = -\frac{1}{3}x + \frac{1}{3}$	CA✓ gradient of perpendicular bisector CA✓ substitution into formula CA ✓ answer (3)

4.1.6 W (-5; 2) into $x + 3y - 1 = 0$ $\begin{aligned}\text{LHS} &= (2(-5) + 6(2) - 2) \\ &= -10 + 12 - 2 \\ &= 0 \\ &= \text{RHS}\end{aligned}$ W is on the line that bisects GZ perpendicularly and W on GZ. $\therefore$ lines intersect at W. OR $\begin{aligned}-\frac{1}{3}x + \frac{1}{3} &= 2 \\ -x + 1 &= 6 \\ x &= -5\end{aligned}$ This is the x - value of the coordinate of W. OR Equation of WZ: $\begin{aligned}y - y_1 &= m(x - x_1) \\ y + 1 &= \frac{2+1}{-5+1}(x+1) \\ y + 1 &= -\frac{3}{4}(x+1) \\ y &= -\frac{3}{4}x - \frac{7}{4} \\ \therefore -\frac{3}{4}x - \frac{7}{4} &= -\frac{1}{3}x + \frac{1}{3} \\ -9x - 21 &= -4x + 4 \\ -5x &= 25 \\ x &= -5 \\ \therefore y &= -\frac{1}{3}(-5) + \frac{1}{3} = 2\end{aligned}$ This is the coordinate of W.	A✓ substitution A✓ = 0 (2) A✓ equating eq. of perpendicular bisector to the horizontal line $y = 2$ A✓ $x = -5$ (2) A✓ equation of WZ A✓ $x = -5$ (2)
4.2.1 circle M: M(-2; 1) ; $r_1 = 5$ circle N: N(1;3) ; $r_2 = 3$ $\therefore r_1 + r_2 = 8$ and $r_1 - r_2 = 2$ $\begin{aligned}\text{MN} &= \sqrt{(1-(-2))^2 + (3-1)^2} \\ &= \sqrt{3^2 + 2^2} \\ &= \sqrt{9 + 4} \\ &= \sqrt{13} \text{ or } 3,6 \\ \therefore r_1 + r_2 &> \text{MN} > r_1 - r_2 \\ \therefore \text{The two circles intersect at two different points.}\end{aligned}$	A ✓ $r_1 = 5$ A ✓ $r_2 = 3$ A ✓ $r_1 + r_2 = 8$ A ✓ $\text{MN} = \sqrt{13}$ = 3,6 A✓ comparing A✓ conclusion (6)


4.2.2 circle M = circle N $(x + 2)^2 + (y - 1)^2 - 25 = (x - 1)^2 + (y - 3)^2 - 9$ $x^2 + 4x + 4 + y^2 - 2y + 1 - 25 = x^2 - 2x + 1 + y^2 - 6y + 9 - 9$ $6x + 4y = 21$ ∴ The equation of the common chord is : $6x + 4y = 21$	M✓ equating A✓ simplifying CA✓ equation of the chord (3) [23]
QUESTION 5 5.1.1 $\tan \alpha = \frac{-2}{2\sqrt{3}} = \frac{-1}{\sqrt{3}}$ ∴ $\alpha = 360^\circ - 30^\circ$ $= 330^\circ$ ∴ $\beta = 30^\circ$ or $\tan(-\beta) = -\tan \beta$ $= -\left(\frac{-2}{2\sqrt{3}}\right)$ $\tan \beta = \frac{2}{2\sqrt{3}}$ $\beta = 30^\circ$	A✓ correct ratio CA✓ $\alpha = 330^\circ$ CA✓ $\beta = 30^\circ$ A✓ correct ratio CACA✓✓ $\beta = 30^\circ$ (3)
5.1.2 $OP^2 = (2\sqrt{3})^2 + (-2)^2 = 12 + 4$ $= 16$ ∴ $OP = 4$	A✓ using distance for formula CA✓ answer (2)
5.1.3 $\frac{OP}{OQ} = \cos \beta$ ∴ $OQ = \frac{OP}{\cos \beta} = \frac{4}{\cos 30^\circ}$ $= \frac{4}{\frac{\sqrt{3}}{2}}$ $= \frac{8}{\sqrt{3}}$	CA✓ $\cos 30^\circ = \frac{\sqrt{3}}{2}$ CA✓ $\frac{8}{\sqrt{3}}$

Downloaded from eduresourceza.com

$\cos \alpha + \sqrt{3} \sin \alpha = k \sin(\alpha + \beta)$ $\cos \alpha + \sqrt{3} \sin \alpha = k(\sin \alpha \cos \beta + \cos \alpha \sin \beta)$ $\cos \alpha + \sqrt{3} \sin \alpha = k \sin \alpha \cos \beta + k \cos \alpha \sin \beta$ $\cos \alpha + \sqrt{3} \sin \alpha = (k \sin \beta) \cos \alpha + (k \cos \beta) \sin \alpha$ $\therefore 1 = k \sin \beta \quad \text{and} \quad \sqrt{3} = k \cos \beta$ $\therefore \sin \beta = \frac{1}{k} \quad \cos \beta = \frac{\sqrt{3}}{k}$   $\sqrt{k^2 - 1} = \sqrt{3}$ $k^2 = 4$ $k = 2$ $\sin \beta = \frac{1}{2}$ $\beta = 30^\circ$	A✓ expansion A✓ comparing coefficients A✓ values of trig ratios A✓ calculating k A✓ calculating β (5) [13]
--	---

QUESTION 6

6.1



A✓asymptotes
 A✓shape of f
 A✓shape of g
 A✓correct x - intercept
 of f (-90° ; 90° ; 270°)
 A✓correct x – intercept
 of g (0° ; 360°)

(5)

6.2 360° A✓ 360°

(1)

6.3 $(0; 3)$ and $(180^\circ; -3)$
 $(-180^\circ; -3)$ and $(360^\circ; 3)$

CA✓for any two

(2)

6.4 $-180^\circ < x < 0^\circ$ or $180^\circ < x < 360^\circ$
 OR
 $-180^\circ < x < 0^\circ \cup 180^\circ < x < 360^\circ$

CA✓ $-180^\circ < x < 0^\circ$
 CA✓ $180^\circ < x < 360^\circ$
 [penalize one mark for
 incorrect notation]

(2)

6.5 $y = 3\cos(x - 45^\circ)$ A✓ $3\cos(x - 45^\circ)$

(1)

[11]


QUESTION 7 7.1 $\cos 54^\circ \cdot \cos x + \sin 54^\circ \cdot \sin x = \sin 2x$ $\cos(54^\circ - x) = \sin 2x$ $= \cos(90^\circ - 2x)$ $\therefore 54^\circ - x = 90^\circ - 2x + n \cdot 360^\circ \text{ or } 54^\circ - x = 360 - (90^\circ - 2x) + n \cdot 360^\circ$ $x = 36^\circ + n \cdot 360^\circ \text{ or } -3x = 216^\circ + n \cdot 360^\circ$ $x = 36^\circ + n \cdot 360^\circ \text{ or } -x = -72^\circ + n \cdot 120^\circ$ $\therefore x = 36^\circ + n \cdot 360^\circ \text{ or } x = -72^\circ + n \cdot 120^\circ, n \in \mathbb{Z}$	A✓ $\cos(54^\circ - x)$ A✓ $\cos(90^\circ - 2x)$ CA✓ $x = 36^\circ + n \cdot 360^\circ$ CA✓ $x = -72^\circ + n \cdot 120^\circ$ A✓ $n \in \mathbb{Z}$ (5)
7.2.1 $\hat{EFD} = \hat{FBC} = \alpha \dots$ corresponding $\angle$'s ... $AD \parallel BC$ $\hat{EFD} = \theta + \hat{AEF} \dots$ ext $\angle$ of $\triangle AEF$ $\therefore \hat{AEF} = \alpha - \theta$ In $\triangle ABE$: $\frac{\sin \hat{AEB}}{AB} = \frac{\sin \hat{EAB}}{BE}$ $\therefore BE = \frac{AB \sin(90^\circ + \theta)}{\sin(\alpha - \theta)}$ $= \frac{AB \cos \theta}{\sin(\alpha - \theta)}$	A✓ S/R A✓ S/R A✓ $\hat{AEF} = \alpha - \theta$ A✓ sine rule application A✓ substitution (5)
7.2.2 Area of $\triangle BCE = \frac{1}{2} \cdot x(18 - 3x) \sin 150^\circ$ $A(x) = \frac{1}{2} x \cdot (18 - 3x) \cdot \frac{1}{2}$ $= \frac{1}{4} \cdot x \cdot 18 - \frac{1}{4} x \cdot 3x$ $= \frac{18}{4} x - \frac{3}{4} x^2$ $= \frac{9}{2} x - \frac{3}{4} x^2$	A✓ area rule A✓ $\sin 150^\circ = \frac{1}{2}$ A✓ simplifying (3)

7.2.3 At maximum area: $A'(x) = 0$ $A'(x) = \frac{9}{2} - 2 \cdot \frac{3}{4}x$ $0 = \frac{9}{2} - \frac{3}{2}x$ $3x = 9$ $x = 3$	M✓ $A'(x) = 0$ A✓ derivative CA✓ $x = 3$ (3)
7.2.4 BC = 3 CE = 18 - 3 (3) = 18 - 9 = 9 BE² = BC² + CE² - 2 BC . CE cos 150° = (3)² + (9)² - 2 x 3 x 9 (-cos 30°) = 9 + 81 + 54 cos 30° = 90 + 54 . $\left(\frac{\sqrt{3}}{2}\right)$ = 136,765 ∴ BE = 11,695 = 11,69	A✓ for both BC = 3 and CE = 9 CA✓ applying cosine rule and substitution CA✓ answer (3) [19]

QUESTION 8

8.1 PT = TQ = 12cm ...(line from center perpendicular to chord PQ) $\therefore PQ = 12 \text{ cm} + 12 \text{ cm} = 24\text{cm}$	A✓ R A✓ answer (2)
8.2 $OT^2 = OQ^2 - QT^2$ pythagoras $= 13^2 - 12^2$ $= 169 - 144$ $= 25$ $\therefore OT = 5$ $\therefore TR = OR - OT$ $= 13\text{cm} - 5\text{cm}$ $= 8\text{cm}$ In ΔPTR, $PR^2 = TR^2 + PT^2$ $= 8^2 + 12^2$ $= 64 + 144$ $= 208 \text{ cm}^2$ $\therefore PR = \sqrt{208} \text{ cm or } 4\sqrt{13} \text{ cm or } 14,42 \text{ cm}$	A✓ OT = 5 CA✓ TR = 8cm CA✓ $PR^2 = 208$ CA✓ $PR = 4\sqrt{13}$ or 14,42 (4) [6]

QUESTION 9

9.1	Interior opposite angle	A✓ S	(1)														
9.2	 Construction : Draw diameter LOQ and join QP or Join OL and OP <table> <tr> <th>STATEMENT</th> <th>REASON</th> </tr> <tr> <td>Let $\hat{P}LM = \hat{L}_1 = x$</td> <td></td> </tr> <tr> <td>$\hat{P}_1 + \hat{P}_2 = 90^\circ$</td> <td>angle subtended by the diameter</td> </tr> <tr> <td>$\hat{L}_2 = 90^\circ - x$</td> <td>$LM \perp OL$, tan – radius</td> </tr> <tr> <td>$\therefore \hat{Q} = x$</td> <td>Sum of the angles of a triangle</td> </tr> <tr> <td>$\hat{N} = x$</td> <td>Subtended by the same chord LP</td> </tr> <tr> <td>$\hat{P}LM = \hat{N}$</td> <td></td> </tr> </table>	STATEMENT	REASON	Let $\hat{P}LM = \hat{L}_1 = x$		$\hat{P}_1 + \hat{P}_2 = 90^\circ$	angle subtended by the diameter	$\hat{L}_2 = 90^\circ - x$	$LM \perp OL$, tan – radius	$\therefore \hat{Q} = x$	Sum of the angles of a triangle	$\hat{N} = x$	Subtended by the same chord LP	$\hat{P}LM = \hat{N}$		A✓ construction A✓ S/R A✓ S/R A✓ S A✓ S/R	(5)
STATEMENT	REASON																
Let $\hat{P}LM = \hat{L}_1 = x$																	
$\hat{P}_1 + \hat{P}_2 = 90^\circ$	angle subtended by the diameter																
$\hat{L}_2 = 90^\circ - x$	$LM \perp OL$, tan – radius																
$\therefore \hat{Q} = x$	Sum of the angles of a triangle																
$\hat{N} = x$	Subtended by the same chord LP																
$\hat{P}LM = \hat{N}$																	
9.3.1	$\hat{A} = 180^\circ - \hat{A}ED \dots$ co interior $\angle$'s, $AB//ED$ $= 180^\circ - 70^\circ$ $= 110^\circ$	A✓ S/R A✓ 110°	(2)														
9.3.2	$\hat{B}_1 = 70^\circ \dots$ ext $\angle$ cyclic quad ABDE	A✓ R A✓ 70°	(2)														
9.3.3	$\hat{D}_2 = \hat{B}_1 = 70^\circ \dots$ (alt $\angle$ s ; $DE//CA$)	CA✓ 70° A✓ S/R	(2)														
9.3.4	$\hat{B}_2 = \hat{D}_2 = 70^\circ \dots$ ($\angle$ s opp = sides)	CA✓ 70° A✓ S/R	(2)														

9.3.5 $\hat{E}_1 = 180^\circ - (\hat{B}_2 + \hat{D}_2) \dots (\angle \text{sum of } \Delta)$ $= 180^\circ - 140^\circ$ $= 40^\circ$ $\therefore \hat{D}_1 = \hat{E}_1 = 40^\circ \dots \text{tan chord theorem}$	CA✓ $\hat{E}_1 = 40^\circ$ CA✓ $\hat{D}_1 = 40^\circ$ A✓ R (3) [17]
--	--

QUESTION 10

10.1 $\hat{P}_1 = \hat{B}_2 = x \dots \text{alt } \angle\text{s}; \text{SP//BC}$ $\hat{P}_2 = \hat{P}_1 = x \dots \text{given}$ $Q_1 = P_1 = x \dots \text{tan chord theorem}$	A✓ S A✓ R A✓ S/R A✓ S/R (4)
10.2 $PC = BC \dots \hat{P}_2 = \hat{B}_2 = x \dots \text{proved above}$ (ΔPCB)	A✓ $\hat{P}_2 = \hat{B}_2 = C = x$ A✓ reason (2)
10.3 $\hat{Q}_1 = \hat{B}_2 = x \dots \text{proved}$ $\therefore \text{RCQB is a cyclic quad}$ $\dots \text{converse } \angle\text{'s in the same segment}$	A✓ S A✓ R (2)
10.4 $\hat{S} = \hat{B}_3 \dots \text{corresp } \angle\text{'s SP} \parallel \text{BC}$ $= \hat{R}_3 \dots \angle\text{'s in the same segment, cyclic quad RCQB}$ In ΔPBS and ΔQCR $\hat{P}_1 = \hat{Q}_1 = x \dots \text{proved}$ $\hat{S} = \hat{R}_3 \dots \text{proved}$ Remaining $\angle\text{s equal}$ $\therefore \Delta PBS \parallel \Delta QCR$	A✓ S/R A✓ S/R A✓ S/R A✓ S/R A✓ R (5)
10.5 In ΔPBQ and ΔPCR $\hat{P}_2$ is common $\hat{PQB} = \hat{R}_2 \dots \text{ext } \angle \text{ of cyclic quad RCQB}$ $\Delta PBQ \parallel \Delta PCR \dots (3^{\text{rd}} \angle \Delta)$ $\therefore \frac{PB}{CP} = \frac{QB}{CR} (\parallel \Delta \text{s})$ $\therefore PB \cdot CR = QB \cdot CP$	A✓ S A✓ S/R A✓ S/R A✓ S (4) [17]

QUESTION 11

In $\triangle KLM$ $\frac{LD}{9} = \frac{8}{6} \dots \quad (\text{LM} \parallel \text{DE}; \text{proportionality theorem})$ $\therefore LD = 12$ $\widehat{DML} = \widehat{MDE} = x \dots \text{alt } \angle s, \text{LM} \parallel \text{DE}$ $\therefore LM = LD = 12 \dots \quad (\text{sides opp } \angle s)$	A✓S/R A✓LD = 12 A✓S A✓answer A✓R (5)
	[5]

TOTAL: [150]