




eduresourceza.com



KWAZULU-NATAL PROVINCE
EDUCATION
REPUBLIC OF SOUTH AFRICA



**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

**MATHEMATICS P1
PREPARATORY EXAMINATION
SEPTEMBER 2022**

MARKS: 150

TIME: 3 hours

This question paper consists of 9 pages and 1 information sheet.

INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 13 questions.
2. Read the questions carefully.
3. Answer ALL the questions.
4. Number your answers exactly as the questions are numbered.
5. Clearly show ALL calculations, diagrams, graphs, etc. which you have used in determining your answers.
6. Answers only will NOT necessarily be awarded full marks.
7. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
8. If necessary, round off answers correct to TWO decimal places, unless stated otherwise.
9. Diagrams are NOT necessarily drawn to scale.
10. Write neatly and legibly.

QUESTION 1

1.1 Solve for x :

$$1.1.1 \quad (x+5)(2x-1)=0 \quad (2)$$

$$1.1.2 \quad -3x^2 - 7x = -8 \quad (\text{correct to TWO decimal places}) \quad (4)$$

$$1.1.3 \quad \sqrt{x+5} + 1 = x \quad (5)$$

$$1.1.4 \quad (2x-3)(x+5) \leq 0 \quad (3)$$

1.2 Solve for x and y simultaneously if:

$$x+3y=5 \text{ and } xy+y^2-3=0 \quad (6)$$

1.3 Simplify fully, without the use of a calculator:

$$\sqrt[n]{\frac{10^n + 2^{n+2}}{5^{2n} + 4 \cdot (5^n)}} \text{ where } n \neq 0 \quad (4)$$

[24]

QUESTION 2

Given the quadratic number pattern: $5 ; 9 ; 17 ; 29 ; \dots$ 2.1 Write down the 5th and 6th terms of the pattern. (2)2.2 Show that the n^{th} term of the quadratic pattern is given by $T_n = 2n^2 - 2n + 5$ (4)2.3 Is 2023 a term in the pattern? Motivate your answer with relevant calculations. (4)
[10]

QUESTION 3

Evaluate: $\sum_{k=1}^{50} (30 - 4k)$

[4]

QUESTION 4

- 4.1 Given the geometric series $a + ar + ar^2 + ar^3 + \dots$, where a is the first term and r is the common ratio. Prove that the sum to n terms of this series is given by

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1 \quad (4)$$

- 4.2 The first two terms of a geometric sequence with constant ratio r , and an arithmetic sequence with constant first difference d , is the same. The first term is 12.


- 4.2.1 Write down the second and third terms of **EACH** sequence in terms of d and r . (2)

- 4.2.2 If it is further given that the sum of the first three terms of the geometric sequence is three more than the sum of the first three terms of the arithmetic sequence. Determine two possible values of the common ratio, r , of the geometric sequence. (5)

[7]

QUESTION 5

Sketched below is the graph of $h(x) = 1 - \frac{1}{x-2}$. A is y-intercept and B is the x-intercept of h .



- 5.1 Write down the equations of the asymptotes of h . (2)
- 5.2 Calculate the coordinates of A and B. (3)
- 5.3 Write down the equation of the line of symmetry of h with positive gradient. (2)
- 5.4 Write down the range of h . (1)


[8]

QUESTION 6

$S(-2; 0)$ and $T(6; 0)$ are the x -intercepts of the graph of $f(x) = ax^2 + bx + c$; $a \neq 0$.

R is the y -intercept of f and g .

The straight line through R and T has the equation $g(x) = -2x + d$.




- 6.1 Calculate the value of d . (2)
- 6.2 Show that $f(x) = -x^2 + 4x + 12$. (4)
- 6.3 Calculate the coordinates of the turning point of f . (3)
- 6.4 Determine for which values of x will:
- 6.4.1 $f(x) - g(x) \geq 0$ (2)
- 6.4.2 $x \cdot f(x) < 0$ (3)
- 6.5 Determine the coordinates of R' , the image of R on p if $p(x) = -f(x-2)$. (2)

[16]

QUESTION 7

Sketched below is the graph of $p(x) = a^x$; $a > 0$; $a \neq 1$. The graph intersects the y -axis at A. The point T (5 ; 32) lies on p .



- 7.1 Write down the coordinates of the point A. (2)
 - 7.2 Calculate the value of a . (2)
 - 7.3 Write down the domain of p . (1)
 - 7.4 Write down the equation of p^{-1} , the inverse of p , in the form $y = \dots$ (2)
 - 7.5 Determine the values of x if $p^{-1}(x) \leq 5$. (2)
- [9]**

QUESTION 8

- 8.1 Dipinda opened an account with an amount of R5000 on 1 June 2021. She then makes monthly deposits of R600 at the end of every month. Her first deposit is made on the 30 June 2021 and her last deposit on 30 April 2023. The account earns interest of 14,25% per annum compounded monthly. Calculate the amount that is in the account directly after her last deposit is made into the account. (6)

- 8.2 Molly wants to buy a house for her family for R800 000. She agreed to pay monthly instalments of R10 000 on the loan which incurred interest at a rate of 13,35 % per annum compounded monthly. The first payment was made at the end of the first month after the loan was granted.

- 8.2.1 Show that the loan will be paid back in full after in 200 months. (4)

- 8.2.2 Suppose Molly encountered unexpected expenses and was unable to pay any instalment at the end of the 120th, 121st, 122nd and 123rd months. At the end of the 124th month she increased her payment to still pay off the loan in 200 months by making 77 equal monthly payments.

- a) Calculate the balance on the loan after the 119th payment was made. (3)

- b) Calculate the new monthly instalment Molly must pay from the 124th month to settle the loan in 200 months. (4)

[17]

QUESTION 9

- 9.1 Determine $f'(x)$ from first principles if $f(x) = \frac{2}{3x}$. (5)

- 9.2 Determine:

- 9.2.1 $g'(x)$ if $g(x) = (x+7)^3$ (5)

- 9.2.2 $\frac{dy}{dx}$ if $y = \sqrt{x^5} - \frac{4}{9x^2}$ (4)

[14]

QUESTION 10

- 10.1 Given: $f(x) = x^3 - 12x - 16$
- 10.1.1 Calculate the x -intercepts of the graph of f . (5)
- 10.1.2 Determine the coordinates of the turning points of f . (4)
- 10.1.3 Sketch the graph of f , indicating the intercepts with the axes and the coordinates of the turning points. (4)
- 10.1.4 Determine the values of x for which the graph of f is concave up. (2)
- 10.2 Given: $p(x) = -x^3 - 8x$
- Is it possible to draw a tangent with positive gradient to the graph of p ? (3)
- Show all calculations to justify your answer. [18]

QUESTION 11

The lead, L , in metres, of a runner in the comrades' marathon in the last t minutes of the race, where $t \in [0; 75]$ is given by equation:

$$L = 1000 + 6t - \frac{t^2}{4}$$

- 11.1 Determine $\frac{dL}{dt}$. (2)
- 11.2 Calculate the time at which the runner has the greatest lead. (2)
- 11.3 At what rate is the runner's lead decreasing when $t = 60$ minutes? (2)
- [6]



QUESTION 12

Given the word “BRACKET”. The letters of this word are randomly arranged to form new arrangements of the letters.

- 12.1 How many unique arrangements of the letters can be made? (2)
- 12.2 Determine the number of unique arrangements of the letters that are randomly made if the letters R and A must be together. (3)
- [5]



QUESTION 13

- 13.1 Two different events A and B are mutually exclusive.


It is further given that:

- $3P(B) = P(A)$
- $P(A \text{ or } B) = 0,63$

Calculate $P(B)$.

(3)

- 13.2 Two cards from a regular pack of 52 playing cards (shown below), are drawn at random one after the other, without replacement.



Calculate the probability that:

- 13.2.1 both cards are picture cards. (2)
- 13.2.2 at least one of the cards is a picture card. (3)
- [8]

TOTAL: 150

INFORMATION SHEET: MATHEMATICS

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2} \{2a + (n - 1)d\}$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; -1 < r < 1$$

$$F = \frac{x[(1 + i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1 + i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}; \frac{y_1 + y_2}{2}\right)$$

$$m = \tan \theta$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cdot \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2} ab \cdot \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2 \sin \alpha \cdot \cos \alpha$$

$$\bar{x} = \frac{\sum fx}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$



education

Department:

Education

PROVINCE OF KWAZULU-NATAL



**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MATHEMATICS P1

PREPARATORY EXAMINATION

SEPTEMBER 2022

MARKING GUIDELINE

MARKS: 150

TIME: 3 hours

NOTE:

- If a candidate answered a QUESTION TWICE, mark only the FIRST attempt.
- If a candidate crossed out an answer and did not redo it, mark the crossed-out answer.
- Consistent accuracy applies to ALL aspects of the marking guidelines.
- Assuming values/answers in order to solve a problem is unacceptable.


This marking guideline consists of 12 pages.

QUESTION 1

1.1.1	$x = -5 \text{ or } x = \frac{1}{2}$	A✓ -5 A✓ $\frac{1}{2}$	(2)
1.1.2	$-3x^2 - 7x + 8 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-(-7) \pm \sqrt{(-7)^2 - 4(-3)(8)}}{2(-3)}$ $x = -3,17 \text{ or } 0,84$	(Penalize 1 mark if rounding off is incorrect – once here for entire paper) A✓ standard form CA✓ correct substitution into quadratic formula CA✓ CA✓ answers	(4)
1.1.3	$\sqrt{x+5} + 1 = x$ $\sqrt{x+5} = x - 1$ $(\sqrt{x+5})^2 = (x-1)^2$ $x+5 = x^2 - 2x + 1$ $x^2 - 3x - 4 = 0$ $(x+1)(x-4) = 0$ $x = -1 \text{ or } x = 4$ n/a	A✓ isolating surd CA✓ standard form CA✓ factors CA✓ $x \neq -1$ CA✓ $x = 4$	(5)
1.1.4	$-5 \leq x \leq \frac{3}{2}$	A✓ critical value -5 A✓ critical value $\frac{3}{2}$ CA✓ interval notation	(3)

1.2	$x + 3y = 5 \rightarrow (1)$ $xy + y^2 - 3 = 0 \rightarrow (2)$ From (1): $x = 5 - 3y \rightarrow (3)$ Substituting (3) into (2): $y(5 - 3y) + y^2 - 3 = 0$ $-2y^2 + 5y - 3 = 0$ $2y^2 - 5y + 3 = 0$ $(2y - 3)(y - 1) = 0$ $y = \frac{3}{2} \text{ or } y = 1$ $x = \frac{1}{2} \text{ or } x = 2$ OR $x + 3y = 5 \rightarrow (1)$ $xy + y^2 - 3 = 0 \rightarrow (2)$ From (1): $y = \frac{5-x}{3} \rightarrow (3)$ Substituting (3) into (2): $x\left(\frac{5-x}{3}\right) + \left(\frac{5-x}{3}\right)^2 - 3 = 0$ $3x(5-x) + 25 - 10x + x^2 - 27 = 0$ $-2x^2 + 5x - 2 = 0$ $2x^2 - 5x + 2 = 0$ $(2x - 1)(x - 2) = 0$ $x = \frac{1}{2} \text{ or } x = 2$ $y = \frac{3}{2} \text{ or } y = 1$	A✓ making x the subject CA✓ substitution CA✓ standard form CA✓ factors CA✓ y – values CA✓ x – values OR A✓ making y the subject CA✓ substitution CA✓ standard form CA✓ factors CA✓ x – values CA✓ y – values	(6)
1.3	$\sqrt[n]{\frac{10^n + 2^{n+2}}{5^{2n} + 4 \cdot 5^n}}$ $= \sqrt[n]{\frac{2^n \cdot 5^n + 2^n \cdot 2^2}{5^{2n} + 4 \cdot 5^n}}$ $= \sqrt[n]{\frac{2^n(5^n + 4)}{5^n(5^n + 4)}}$ $= \sqrt[n]{\frac{2^n}{5^n}}$ $= \frac{2}{5}$	A✓ factorising numerator A✓ factorising denominator CA✓ simplifying CA✓ answer	(4)
[24]			

QUESTION 2

2.1	45 ; 65	AA✓✓ answers	(2)
2.2	 1D 2D $2a = 4 \quad \therefore a = 2$ $3a + b = 4 \quad \therefore b = -2$ $a + b + c = 5 \quad \therefore c = 5$ $T_n = 2n^2 - 2n + 5$	A✓ $2a = 4$ A✓ $a = 2$ A✓ $3a + b = 4$ A✓ $a + b + c = 5$	(4)
2.3	$T_n = 2n^2 - 2n + 5 = 2023$ $2n^2 - 2n - 2018 = 0$ $n^2 - n - 1009 = 0$ $n = \frac{1 \pm \sqrt{1 + 4036}}{2} = 32.27 \text{ or } -31.27$ Since n is not a Natural Number, 2023 is not a term of the sequence.	A✓ equating n^{th} term to 2023 CA✓ standard form CA✓ n – values CA✓ conclusion	(4)
			[10]

QUESTION 3

26 ; 22 ; 18 ; ... $S_n = \frac{n}{2} [2a + (n - 1)d]$ $S_{50} = \frac{50}{2} [2(26) + (50 - 1)(-4)]$ $S_{50} = -3600$ OR $S_n = \frac{n}{2} [a + T_n]$ $S_{50} = \frac{50}{2} [26 + (-170)]$ $S_{50} = -3600$	A✓ n – value A✓ a – value A✓ d – value CA✓ answer OR A✓ n – value A✓ a – value A✓ T_{50} – value CA✓ answer	(4) (4)	
			[4]

QUESTION 4

4.1	$S_n = a + ar + ar^2 + ar^3 + \dots + ar^{n-1} \rightarrow (1)$ $rS_n = ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n \rightarrow (2)$ $(2) - (1):$ $rS_n - S_n = ar^n - a$ $S_n(r - 1) = a(r^n - 1)$ $S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$	A✓ equation (1) A✓ equation (2) A✓ subtracting LHS and RHS terms A✓ factorising	(4)
4.2.1	AS: $12 + d; 12 + 2d; \dots$ GS: $12r; 12r^2; \dots$	A✓ AS set up of terms A✓ GS set up of terms	(2)
4.2.2	$12 + d = 12r \rightarrow (1)$ $36 + 3d + 3 = 12 + 12r + 12r^2 \rightarrow (2)$ From (1): $d = 12r - 12 \rightarrow (3)$ Substituting (3) into (2), we have $36 + 3(12r - 12) + 3 = 12 + 12r + 12r^2$ $12 + (12r - 12) + 1 = 4 + 4r + 4r^2$ $4r^2 - 8r + 3 = 0$ $(2r - 1)(2r - 3) = 0$ $r = \frac{1}{2} \text{ or } r = \frac{3}{2}$	A✓ equation (1) and (2) A✓ making d the subject CA✓ standard quadratic form CA✓ factors CA✓ answers	(5)
			[11]

QUESTION 5


5.1	$y = 1 - \frac{1}{x - 2}$ $x = 2 \text{ and } y = 1$	A✓ $x = 2$ A✓ $y = 1$	(2)
5.2	y - intercept: $(0; 1\frac{1}{2})$ x - intercept: $x = 3$ $(3; 0)$	A✓ y -intercept A✓ $x = 3$ CA✓ coordinate form	(3)
5.3	$y = x - 1$	A✓ Gradient value A✓ y - intercept	(2)
5.4	$y \in R; y \neq 1$ OR $y \in (-\infty; 1) \cup (1; \infty)$	A✓ answer OR A✓ answer	(1) (1)
			[8]

Downloaded from eduresourceza.com

QUESTION 7

7.1	$A(0; 1)$	AA✓✓ answer	(2)
7.2	$y = a^x$ $32 = a^5$ $a = 2$	A✓ substitution of point T(5;32) A✓ answer	(2)
7.3	$x \in R$ OR $x \in (-\infty; \infty)$	A✓ answer OR A✓ answer	(1) (1)
7.4	$y = \log_2 x$	CACA✓✓ answer	(2)
7.5	$\log_2 x = 5$ $x = 2^5 = 32$ $0 < x \leq 32$	CA✓ end points A✓ interval Can be solved by log inequalities. Answer Only – Full marks	(2)
			[9]

QUESTION 8


8.1	<table border="1"> <tr> <td>1 Jun 2021</td> <td>31 Jul 2021</td> <td>31 Aug 2021</td> <td>.....</td> <td>30 Apr 2023</td> </tr> <tr> <td>5000</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>600</td> <td>600</td> <td>600</td> <td>.....</td> <td>600</td> </tr> </table> Dipinda's final amount in the account: $= P(1+i)^n + \frac{x[(1+i)^n - 1]}{i}$ $= 5000 \left(1 + \frac{14.25\%}{12}\right)^{23} + \frac{600 \left[\left(1 + \frac{14.25\%}{12}\right)^{23} - 1\right]}{\frac{14.25\%}{12}}$ $= R22\,321,54$	1 Jun 2021	31 Jul 2021	31 Aug 2021		30 Apr 2023	5000					600	600	600		600	<u>A – formula</u> A✓ value of n A✓ value of i <u>FV – formula</u> A✓ value of n CA✓ correct substitution into A CA✓ correct substitution into Fv CA✓ answer	(6)
1 Jun 2021	31 Jul 2021	31 Aug 2021		30 Apr 2023														
5000																		
600	600	600		600														
8.2.1	$P = \frac{x[1 - (1+i)^{-n}]}{i}$ $800\,000 = \frac{10000 \left[1 - \left(1 + \frac{13.35\%}{12}\right)^{-n}\right]}{\frac{13.35\%}{12}}$ $\left(1 + \frac{13.35\%}{12}\right)^{-n} = \frac{11}{100} = 0,11$ $-n = \log_{\left(1 + \frac{13.35\%}{12}\right)} 0,11$ $n = 199,5083362$ Therefore the loan will be paid off in 200 months. N.B. Candidates can also substitute the value of 200 into the Pv formula to show that the loan will be paid in 200 months.	 A✓ value of P, x and value of i A✓ substitution into formula A✓ use of logs A✓ decimal value	(4)															

8.2.2a	$P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $= \frac{10\,000 \left[1 - \left(1 + \frac{13.35\%}{12} \right)^{-80,5083362} \right]}{\frac{13.35\%}{12}}$ $= R530\,009,55$ If $n = 81$ is used and $P = R532\,010,58$ Give a maximum of 2/3 marks N.B. Candidates can also use the method of A – Fv	A✓ value of n A✓ value of i CA✓ answer	(3)
8.2.2b	$A = P(1 + i)^n$ $A = R530\,009,55 \left(1 + \frac{13.35\%}{12} \right)^4$ $A = R\,553\,991,4839$ $P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $553\,991,4839 = \frac{x \left[1 - \left(1 + \frac{13.35\%}{12} \right)^{-77} \right]}{\frac{13.35\%}{12}}$ $x = R10\,748,55$	A✓ value of n CA✓ answer A✓ value of n CA✓ answer	(4)
[17]			

QUESTION 9 (penalize 1 mark once for incorrect notation in this question)

9.1	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{\frac{2}{3(x+h)} - \frac{2}{3x}}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{2x - 2(x+h)}{3x(x+h)} \times \frac{1}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{2x - 2x - 2h}{3x(x+h)} \times \frac{1}{h}$ $f'(x) = \frac{-2}{3x^2}$ OR $f(x+h) = \frac{2}{3(x+h)}$ $f(x+h) - f(x) = \frac{2}{3(x+h)} - \frac{2}{3x}$ $\frac{f(x+h) - f(x)}{h} = \frac{-2}{3x(x+h)}$ $f'(x) = \lim_{h \rightarrow 0} \frac{-2}{3x(x+h)}$ $f'(x) = \frac{-2}{3x^2}$	A✓ formula A✓ substitution CA✓ LCD CA✓ simplification of numerator CA✓ answer OR A✓ value of $f(x+h)$ CA✓ value of $f(x+h) - f(x)$ CA✓ value of $\frac{f(x+h) - f(x)}{h}$ A✓ formula CA✓ answer	(5) (5)
9.2.1	$g(x) = (x+7)^3$ $g(x) = x^3 + 21x^2 + 147x + 343$ $g'(x) = 3x^2 + 42x + 147$	AA✓✓ (two terms correct 1 mark, all terms correct 2 marks) CACACA✓✓✓ each term	(5)
9.2.2	$y = \sqrt{x^5} - \frac{4}{9x^2}$ $y = x^{\frac{5}{2}} - \frac{4}{9}x^{-2}$ $\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}} + \frac{8}{9}x^{-3}$	AA✓✓ writing in exponential form CACA✓✓ each term	(4)
[14]			

QUESTION 10

10.1.1	$x^3 - 12x - 16 = 0$ $(x + 2)(x^2 - 2x - 8) = 0$ $(x + 2)(x + 2)(x - 4) = 0$ $x = -2 \text{ or } x = 4$	A✓ binomial factor AA✓✓ factors CA CA ✓✓ answers	(5)
10.1.2	$f(x) = x^3 - 12x - 16$ $f'(x) = 3x^2 - 12 = 0$ $x^2 - 4 = 0$ $(x + 2)(x - 2) = 0$ $x = -2 \text{ or } x = 2$ $y = 0 \text{ or } y = -32$	A✓ derivative and equating to 0 CA✓ factors CA✓ x – values CA✓ y – values	(4)
10.1.3		CA✓ Maximum and Minimum points CA✓ x – intercepts A✓ y – intercept A✓ shape	(4)
10.1.4	$f''(x) = 6x > 0$ $x > 0$	A✓ 2 nd derivative A✓ answer	(2)
10.2	$p'(x) = -3x^2 - 8$ $-3x^2 \leq 0 \text{ for all } x \in \mathbb{R}$ $-3x^2 - 8 \leq 0$ The gradient of all tangents to the graph of p is always negative.	A✓ derivative A✓ reasoning A✓ reasoning	(3)
			[18]

QUESTION 11

11.1	$L = 1000 + 6t - \frac{t^2}{4}$ $\frac{dL}{dt} = 6 - \frac{1}{2}t$	AA✓✓ for each term	(2)
11.2	For greatest lead: $\frac{dL}{dt} = 0$ $6 - \frac{1}{2}t = 0$ $t = 12$ minutes 	CA✓ $\frac{dL}{dt} = 0$ or equating derivative to 0 CA✓ answer	(2)
11.3	$\frac{dL}{dt}_{t=60} = 6 - \frac{1}{2}(60)$ $\frac{dL}{dt}_{t=60} = -24$ The runner's lead is decreasing at 24 metres per minute	CA✓ substitution of $t = 60$ into derivative and value of -24 CA✓ conclusion (provided the derivative is $-ve$)	(2)
			[6]

QUESTION 12

12.1	7! or 5 040	A✓ A✓ 7! or 5040	(2)
12.2	$6! \times 2!$ $= 1440$	AA✓✓ $6! \times 2!$ A✓ 1440	(3)
			[5]

QUESTION 13

13.1	$P(A \text{ or } B) = P(A) + P(B)$ $0,63 = 3P(B)+P(B)$ $4P(B) = 0,63$ $P(B) = 0,16$	A✓ condition for mutually exclusive events A✓ correct substitution A✓ P(B) value	(3)																																										
13.2.1	$P(\text{Both Picture cards}) = \frac{12}{52} \times \frac{11}{51}$ $= \frac{11}{221} = 0,0498 = 4,98 \%$	A✓ $\frac{12}{52} \times \frac{11}{51}$ A ✓ $\frac{11}{221} = 0,0498 = 4,98 \%$	(2)																																										
13.2.2	<table border="1"><thead><tr><th></th><th>1ST</th><th></th><th>2ND</th><th>Outcomes</th><th>Probabilities</th></tr></thead><tbody><tr><td></td><td></td><td>11/51 → P</td><td>P</td><td>PP</td><td>$\frac{12}{52} \times \frac{11}{51}$</td></tr><tr><td></td><td></td><td>40/51 → NP</td><td>NP</td><td>PNP</td><td>$\frac{12}{52} \times \frac{40}{51}$</td></tr><tr><td>12/52 → P</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td></td><td></td><td>12/51 → P</td><td>P</td><td>NPP</td><td>$\frac{40}{52} \times \frac{12}{51}$</td></tr><tr><td></td><td></td><td>39/51 → NP</td><td>NP</td><td>NPNP</td><td>$\frac{40}{52} \times \frac{39}{51}$</td></tr><tr><td>40/52 → NP</td><td></td><td></td><td></td><td></td><td></td></tr></tbody></table> P (at least 1 picture card) $= 1 - P(\text{no picture card})$ $= 1 - \left(\frac{40}{52} \times \frac{39}{51}\right)$ $= \frac{7}{17} = 0,4118 = 41,18 \%$ OR P (at least 1 picture) $= \left(\frac{12}{52} \times \frac{11}{51}\right) + \left(\frac{12}{52} \times \frac{40}{51}\right) + \left(\frac{40}{52} \times \frac{12}{51}\right)$ $= \frac{7}{17} = 0,4118 = 41,18 \%$		1 ST		2 ND	Outcomes	Probabilities			11/51 → P	P	PP	$\frac{12}{52} \times \frac{11}{51}$			40/51 → NP	NP	PNP	$\frac{12}{52} \times \frac{40}{51}$	12/52 → P								12/51 → P	P	NPP	$\frac{40}{52} \times \frac{12}{51}$			39/51 → NP	NP	NPNP	$\frac{40}{52} \times \frac{39}{51}$	40/52 → NP						A✓ Method A✓ Correct Substitution A✓ $\frac{7}{17} = 0,4118 = 41,18 \%$ OR AA✓✓probabilities A✓ $\frac{7}{17} = 0,4118 = 41,18 \%$	(3)
	1 ST		2 ND	Outcomes	Probabilities																																								
		11/51 → P	P	PP	$\frac{12}{52} \times \frac{11}{51}$																																								
		40/51 → NP	NP	PNP	$\frac{12}{52} \times \frac{40}{51}$																																								
12/52 → P																																													
		12/51 → P	P	NPP	$\frac{40}{52} \times \frac{12}{51}$																																								
		39/51 → NP	NP	NPNP	$\frac{40}{52} \times \frac{39}{51}$																																								
40/52 → NP																																													
			[8]																																										