



eduresourceza.com



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE
NASIONALE
SENIOR SERTIFIKAAT**

GRADE/GRAAD 12

MATHEMATICS P1/WISKUNDE V1

NOVEMBER 2013

MEMORANDUM

MARKS/PUNTE: 150

**This memorandum consists of 24 pages.
*Hierdie memorandum bestaan uit 24 bladsye.***

NOTE:

- If a candidate answered a question TWICE, mark only the first attempt.
- If a candidate crossed out an attempt of a question and did not redo the question, mark the crossed-out question.
- Consistent accuracy applies in ALL aspects of the marking memorandum.
- Assuming values/answers in order to solve a problem is unacceptable.




LET WEL:

- As 'n kandidaat 'n vraag TWEE keer beantwoord het, merk slegs die eerste poging.
- As 'n kandidaat 'n antwoord deurgehaal en nie oorgedoen het nie, merk die deurgehaalde antwoord.
- Volgehoue akkuraatheid is DEURGAANS in ALLE aspekte van die memorandum van toepassing.
- Aanvaarding van waardes/antwoorde om 'n problem op te los, is onaanvaarbaar.

QUESTION/VRAAG 1

1.1.1	$x^2 - x - 12 = 0$ $(x - 4)(x + 3) = 0$ $x = 4 \quad \text{or} \quad x = -3$ NOTE: Answer only: max 2/3 marks OR $x^2 - x - 12 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-12)}}{2(1)}$ $= \frac{1 \pm \sqrt{49}}{2}$ $= 4 \quad \text{or} \quad -3$	NOTE: Answer only: max 2/3 marks ✓ factors ✓ answer ✓ answer (3)
1.1.2 (a)	$2x^2 - 5x - 11 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-5) \pm \sqrt{(-5)^2 - 4(2)(-11)}}{2(2)}$ $= \frac{5 \pm \sqrt{113}}{4}$ $= 3,91 \quad \text{or} \quad -1,41$ OR NOTE: • Answer only: max 2/4 marks • If the answer is left as $\frac{5 \pm \sqrt{113}}{4}$: 4/4 marks • If candidate continues after correct surd but then has incorrect answers: max 4/4 marks	NOTE: Wrong formula: 0/4 marks ✓ correct substitution of b into correct formula ✓ correct substitution of a and c into correct formula ✓ ✓ $\frac{5 \pm \sqrt{113}}{4}$ OR decimal answers (4)

	$2x^2 - 5x - 11 = 0$ $x^2 - \frac{5}{2}x = \frac{11}{2}$ $\left(x - \frac{5}{4}\right)^2 = \frac{11}{2} + \frac{25}{16}$ $\left(x - \frac{5}{4}\right) = \pm \sqrt{\frac{113}{16}}$ $x = \frac{5}{4} \pm \sqrt{\frac{113}{16}}$ $x = 3,91 \quad \text{or} \quad x = -1,41$ OR $2x^2 - 5x - 11 = 0$ $x^2 - \frac{5x}{2} - \frac{11}{2} = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-\left(-\frac{5}{2}\right) \pm \sqrt{\left(-\frac{5}{2}\right)^2 - 4(1)\left(-\frac{11}{2}\right)}}{2(1)}$ $= \frac{\frac{5}{2} \pm \sqrt{\frac{113}{4}}}{2}$ $x = 3,91 \quad \text{or} \quad x = -1,41$	✓ division by 2 ✓ $\left(x - \frac{5}{4}\right) = \pm \sqrt{\frac{113}{16}}$ ✓ $x = \frac{5}{4} \pm \sqrt{\frac{113}{16}}$ ✓ answers (4) ✓ division by 2 ✓ subs into correct formula ✓ $\frac{\frac{5}{2} \pm \sqrt{\frac{113}{4}}}{2}$ ✓ answer (4)																														
1.1.2 (b)	$2x^3 - 5x^2 - 11x = 0$ $x(2x^2 - 5x - 11) = 0$ $x = 0 \quad \text{or} \quad x = 3,91 \quad \text{or} \quad x = -1,41$ OR $x = 0 \quad \text{or} \quad x = \frac{5 \pm \sqrt{113}}{4}$ NOTE: • Division by x: max 1/2 marks • Use quadratic formula to solve cubic: 0/2 marks • Answer only : 2/2 marks	✓ factors ✓ answers (2)																														
1.1.3	$-3(x+7)(x-5) < 0$ $\begin{array}{ccccccc} - & 0 & + & 0 & - & & \\ & -7 & & 5 & & & \end{array} \quad \text{OR} \quad \begin{array}{c} \text{Graph of } y = -3(x+7)(x-5) \\ \text{The parabola opens downwards with x-intercepts at } -7 \text{ and } 5. \end{array}$ OR <table><tr><td></td><td></td><td>-7</td><td></td><td>5</td><td></td></tr><tr><td>x</td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>x + 7</td><td></td><td>+</td><td>0</td><td>-</td><td>-</td></tr><tr><td>x - 5</td><td></td><td>+</td><td>+</td><td>0</td><td>-</td></tr><tr><td>-3(x + 7)(x - 5)</td><td></td><td>-</td><td>0</td><td>+</td><td>0</td></tr></table> $x < -7 \quad \text{or} \quad x > 5 \quad \text{OR} \quad x \in (-\infty; -7) \cup (5; \infty)$			-7		5		x						x + 7		+	0	-	-	x - 5		+	+	0	-	-3(x + 7)(x - 5)		-	0	+	0	✓ critical values ✓ $x < -7$ ✓ $x > 5$ ✓ or / ∪
		-7		5																												
x																																
x + 7		+	0	-	-																											
x - 5		+	+	0	-																											
-3(x + 7)(x - 5)		-	0	+	0																											

OR  OR $-3(x+7)(x-5) < 0$ $(x+7)(x-5) > 0$ $\begin{array}{ccccccc} + & 0 & - & 0 & + \\ -7 & & & 5 & \end{array}$ OR  OR <table style="margin-left: auto; margin-right: auto; border-collapse: collapse;"> <tr> <td style="padding: 5px;">x</td> <td style="border-bottom: 1px solid black; padding: 5px;"></td> <td style="border-bottom: 1px solid black; padding: 5px;">-7</td> <td style="border-bottom: 1px solid black; padding: 5px;"></td> <td style="border-bottom: 1px solid black; padding: 5px;">5</td> <td style="border-bottom: 1px solid black; padding: 5px;"></td> </tr> <tr> <td style="padding: 5px;">$x+7$</td> <td style="padding: 5px;">$-$</td> <td style="padding: 5px;">0</td> <td style="padding: 5px;">$+$</td> <td style="padding: 5px;">$+$</td> <td style="padding: 5px;">$+$</td> </tr> <tr> <td style="padding: 5px;">$x-5$</td> <td style="padding: 5px;">$-$</td> <td style="padding: 5px;">$-$</td> <td style="padding: 5px;">$-$</td> <td style="padding: 5px;">0</td> <td style="padding: 5px;">$+$</td> </tr> <tr> <td style="padding: 5px;">$(x+7)(x-5)$</td> <td style="padding: 5px;">$+$</td> <td style="padding: 5px;">0</td> <td style="padding: 5px;">$-$</td> <td style="padding: 5px;">0</td> <td style="padding: 5px;">$+$</td> </tr> </table> $x < -7$ or $x > 5$ OR $x \in (-\infty; -7) \cup (5; \infty)$ OR  NOTE: In this alternative, award max 3/4 marks since there is no conclusion	x		-7		5		$x+7$	$-$	0	$+$	$+$	$+$	$x-5$	$-$	$-$	$-$	0	$+$	$(x+7)(x-5)$	$+$	0	$-$	0	$+$	(4) ✓critical values
x		-7		5																					
$x+7$	$-$	0	$+$	$+$	$+$																				
$x-5$	$-$	$-$	$-$	0	$+$																				
$(x+7)(x-5)$	$+$	0	$-$	0	$+$																				

1.2	$y + 2 = x \text{ and } y = x^2 - x - 10$ $y + 2 = x$ $y = (y + 2)^2 - (y + 2) - 10$ $y = y^2 + 4y + 4 - y - 2 - 10$ $0 = y^2 + 2y - 8$ $0 = (y + 4)(y - 2)$ $y = -4 \text{ or } 2$ $x = -4 + 2 \text{ or } x = 2 + 2$ $= -2 \qquad \qquad = 4$ OR $y + 2 = x \text{ and } y = x^2 - x - 10$ $x^2 - x - 10 + 2 = x$ $0 = x^2 - 2x - 8$ $0 = x^2 - 2x - 8$ $0 = (x - 4)(x + 2)$ $x = 4 \text{ or } -2$ $y = 4 - 2 \text{ or } y = -2 - 2$ $= 2 \qquad \qquad = -4$ OR $y + 2 = x \text{ and } y = x^2 - x - 10$ $y = x - 2$ $x - 2 = x^2 - x - 10$ $0 = x^2 - 2x - 8$ $0 = x^2 - 2x - 8$ $0 = (x - 4)(x + 2)$ $x = 4 \text{ or } -2$ $y = 4 - 2 \text{ or } y = -2 - 2$ $= 2 \qquad \qquad = -4$	✓ substitution ✓ $0 = y^2 + 2y - 8$ ✓ factors ✓ y-values ✓✓ x-values (6) ✓ substitution ✓ $0 = x^2 - 2x - 8$ ✓ factors ✓ x-values ✓✓ y-values (6) ✓ substitution ✓ $0 = x^2 - 2x - 8$ ✓ factors ✓ x-values ✓✓ y-values (6)
1.3	$\frac{3^{2015} + 3^{2013}}{9^{1006}}$ $= \frac{3^{2013}(3^2 + 1)}{3^{2012}}$ $= 3(10)$ $= 30$ OR	✓ $3^{2013}(3^2 + 1)$ ✓ denominator ✓ answer (3)

$\frac{3^{2015} + 3^{2013}}{9^{1006}}$ $= \frac{3^{2012}(3^3 + 3)}{3^{2012}}$ $= 27 + 3$ $= 30$ OR Let $x = 3^{2012}$ $\frac{3^{2015} + 3^{2013}}{9^{1006}}$ $= \frac{3^{2012} \cdot 3^3 + 3^{2012} \cdot 3}{3^{2012}}$ $= \frac{27x + 3x}{x}$ $= \frac{30x}{x}$ $= 30$	$\checkmark 3^{2012}(3^3 + 3)$ $\checkmark \text{ denominator}$ $\checkmark \text{ answer}$ (3) $\checkmark 3^{2012} \cdot 3^3 + 3^{2012} \cdot 3 / 27x + 3x$ $\checkmark \text{ denominator}$ $\checkmark \text{ answer}$ (3) [22]
---	---

QUESTION/VRAAG 2


2.1	Given geometric sequence : 7 ; x ; 63 ... $\frac{T_2}{T_1} = \frac{T_3}{T_2}$ $\frac{x}{7} = \frac{63}{x}$ $x^2 = 441$ $x = \pm 21$ OR Given geometric sequence : 7 ; x ; 63 ... $\frac{T_2}{T_1} = \frac{T_3}{T_2}$ $\frac{x}{7} = \frac{63}{x}$ $x^2 = 441$ $x^2 - 441 = 0$ $(x - 21)(x + 21) = 0$ $x = \pm 21$ OR $63 = 7r^2$ $r^2 = 9$ $r = \pm 3$ $x = \pm 21$	$\checkmark \frac{T_2}{T_1} = \frac{T_3}{T_2} / \frac{x}{7} = \frac{63}{x}$ $\checkmark x^2 = 441$ $\checkmark \text{both answers}$ (3) $\checkmark \frac{T_2}{T_1} = \frac{T_3}{T_2} / \frac{x}{7} = \frac{63}{x}$ $\checkmark x^2 = 441$ $\checkmark \text{both answers}$ (3) $\checkmark 63 = 7r^2$ $\checkmark r^2 = 9$ $\checkmark \text{both answers}$ (3)
2.2.1	$r = \frac{10}{15} = \frac{2}{3}$ $T_n = ar^{n-1}$ $T_{10} = 15 \left(\frac{2}{3} \right)^{10-1}$ $= \frac{2560}{6561} \text{ or } 0,39$ OR $r = \frac{10}{15} = \frac{2}{3}$ Expansion of the series $15 + 10 + \frac{20}{3} + \frac{40}{9} + \frac{80}{27} + \frac{160}{81} + \frac{320}{243} + \frac{640}{729} + \frac{1280}{2187} + \frac{2560}{6561}$ $T_{10} = \frac{2560}{6561}$ NOTE: If the candidate rounds off early and gets $r = 0,67$, then $T_{10} = 0,41$: 3/3 marks	$\checkmark r = \frac{2}{3}$ $\checkmark \text{correct subs into correct formula}$ $\checkmark \text{answer}$ (3) $\checkmark r = \frac{2}{3}$ $\checkmark \text{expansion of the series}$ $\checkmark \text{answer}$ (3)

2.2.2	$S_n = \frac{a(r^n - 1)}{r - 1}$ $S_9 = \frac{15\left(\left(\frac{2}{3}\right)^9 - 1\right)}{\frac{2}{3} - 1}$ $= \frac{95855}{2187}$ $= 43,83$ OR $S_n = \frac{a(1 - r^n)}{1 - r}$ $S_9 = \frac{15\left(1 - \left(\frac{2}{3}\right)^9\right)}{1 - \left(\frac{2}{3}\right)}$ $= \frac{95855}{2187}$ $= 43,83$	✓ correct substitution into correct formula ✓ answer (2)
2.3.1	$T_{191} = 0$	✓ answer (1)
2.3.2	Since the sum of all odd-positioned terms will be zero, need only consider the sum of the even-positioned terms, which form an arithmetic sequence, i.e. the sum of 250 even terms: <i>Omdat die som van al die terme in onewe posisies nul is, slegs nodig om die som van die terme in ewe posisies te oorweeg, wat 'n rekenkundige ry vorm, m.a.w. die som van 250 ewe terme:</i> $S_{500} = \frac{250}{2} \left[2\left(-\frac{1}{2}\right) + (250-1)(1) \right]$ $= 31000$ NOTE: Breakdown: If $n = 500$ with $a = -\frac{1}{2}$ and $d = 1$ then $S_n = 124\,500$: max 2/4 marks OR $S_{500} = \frac{125[2(-1) + 249(2)]}{2}$ $= 31000$ OR	✓ $n = 250$ ✓ $a = -\frac{1}{2}$ and $d = 1$ ✓ substitution into correct formula ✓ answer ✓ $n = 125$ ✓ $a = -1$ and $d = 2$ ✓ subs into correct formula ✓ answer (4)

	$\frac{3}{2} + \frac{5}{2} + \dots \text{to } 248 \text{ terms}$ $= 124 \left[\frac{3}{2} + \frac{497}{2} \right]$ $= 124 \times 250$ $= 31000$ OR $\frac{3}{2} + \frac{5}{2} + \dots \text{to } 248 \text{ terms}$ $= 124[3 + 247]$ $= 124 \times 250$ $= 31000$ OR Sum = $0 + 4 + 8 + \dots$ to 125 terms $= \frac{125}{2} [0 + (125 - 1)4]$ $= 31000$	✓ $n = 248$ ✓ subs into correct formula ✓ $\frac{3}{2} + \frac{247}{2}$ ✓ answer (4) ✓ $n = 248$ ✓ subs into correct formula ✓ $3 + 247$ ✓ answer (4) ✓ $n = 125$ ✓ $a = 0$ and $d = 4$ ✓ subs into correct formula ✓ answer (4)
2.4.1	$T_1 = (4(1) - 1)^2$ $= 3^2$ $= 9$ NOTE: If $k = 1$, $T_1 = 3$: max 1/2 mark	✓ subs $x = 1$ and $k = 2$ ✓ answer (2)
2.4.2	$r = 4x - 1$ $-1 < r < 1$ $-1 < 4x - 1 < 1$ $0 < 4x < 2$ $0 < x < \frac{1}{2}$ NOTE: Incorrect r: max 1/3 marks If candidate only writes down $4x - 1$ and does nothing else: 0/3 marks	✓ $r = 4x - 1$ ✓ $-1 < 4x - 1 < 1$ ✓ answer (3)

[18]

QUESTION/VRAAG 3


3.1.1	$T_n = 4n - 7$ OR $T_n = -3 + (n-1)(4)$	$\checkmark 4n$ $\checkmark -7$ $\checkmark -3$ $\checkmark (n-1)(4)$ (2)
3.1.2	$T_4 = 9$ $T_5 = 13$ $T_6 = 17$ $T_7 = 21$	$\checkmark$ any TWO consecutive answers correct $\checkmark$ last TWO answers correct (2)
3.1.3	0 ; 1 ; 2 ; 0 ; 1 ; 2 ; 0	2 marks for all 7 correct OR 1 mark for only first / last 3 correct OR 0 marks if less than 3 correct (2)
3.1.4	Multiples of 3 in the pattern are: $-3 ; 9 ; 21$ $T_n = -3 + 12(n-1)$ $T_n = a + (n-1)d$ $T_n = 12n - 15$ $393 = -3 + (n-1)(12)$ $393 = 12n - 15$ or $393 = 12n - 15$ $12n = 408$ $12n = 408$ $n = 34$ $n = 34$ $S_n = \frac{n}{2}[a + L]$ $S_n = \frac{n}{2}[2a + (n-1)d]$ $S_{34} = \frac{34}{2}[-3 + 393]$ or $S_{34} = \frac{34}{2}[2(-3) + 33(12)]$ $S_{34} = 6630$ $S_{34} = 6630$ NOTE: • If the candidate does not show the working to get to $n = 34$: no penalty • If a candidate sums the whole sequence: 0/5 marks • Answer only: max 1/5 marks	$\checkmark 12n - 15$ $\checkmark 393 = 12n - 15$ $\checkmark n = 34$ $\checkmark$ subs $a = -3$ and $d = 12$ into correct formula $\checkmark S_{34} = 6630$ (5)
3.2.1	 $T_5 = 35$	$\checkmark \checkmark$ answer (2)

	OR The sequence is 1, 5, 12, 22, 35. Therefore $T_5 = 35$ OR $T_5 = 22 + 13 = 35$	✓✓ answer (2) ✓✓ answer (2)
3.2.2	$T_{50} = T_1 + \frac{49}{2}[2(4) + 48(3)]$ $= 1 + 3724$ $= 3725$ OR $2a = 3$ $a = \frac{3}{2}$ $3\left(\frac{3}{2}\right) + b = 4$ $b = -\frac{1}{2}$ $\left(\frac{3}{2}\right) + \left(-\frac{1}{2}\right) + c = 1$ $c = 0$ $T_n = \frac{3}{2}n^2 - \frac{1}{2}n$ $T_{50} = \frac{3}{2}(50)^2 - \frac{1}{2}(50)$ $= 3725$ OR $T_1 = 1$ $T_2 - T_1 = 4$ $T_3 - T_2 = 7$ $T_4 - T_3 = 10$... $T_{50} - T_{49} = ?$ Add both sides $T_{50} = 1 + 4 + 7 + 10 + \dots$ to 50 terms $= \frac{50}{2}(2 + 49(3))$ $= 3725$ NOTE: • Answer only: max 1 mark • If the candidate calculates the general formula in 3.2.1, they can be awarded 5/5 marks in 3.2.2	✓ $a = 4$ ✓ $d = 3$ ✓ $n = 49$ ✓ substitution into correct formula ✓ answer (5) ✓ $a = \frac{3}{2}$ ✓ $b = -\frac{1}{2}$ ✓ $c = 0$ ✓ subs $n = 50$ ✓ answer (5) ✓✓ expansion ✓ $T_{50} = 1 + 4 + 7 + 10 + \dots$ to 50 terms ✓ subs into correct formula ✓ answer (5)


QUESTION/VRAAG 4


Given: $f(x) = -2x^2 - 5x + 3$

4.1	$(0 ; 3)$ OR $x = 0$ and $y = 3$	- If there is evidence that the candidate has indicated that $x = 0$ and $y = 3$: 1 mark - If candidate just states $y = 3$: 1 mark	✓ answer in coordinate form (1) ✓ both values correct (1)
4.2	$0 = -2x^2 - 5x + 3$ $0 = 2x^2 + 5x - 3$ $0 = (2x - 1)(x + 3)$ $x = \frac{1}{2}$ or -3		✓ $y = 0$ ✓ factors ✓ x -values (3)
4.3	$f(x) = -2x^2 - 5x + 3$ $x = -\frac{b}{2a}$ $= -\frac{(-5)}{2(-2)}$ $= -\frac{5}{4}$ or $-1\frac{1}{4}$ $y = f\left(-\frac{5}{4}\right)$ $= -2\left(-\frac{5}{4}\right)^2 - 5\left(-\frac{5}{4}\right) + 3$ $= \frac{49}{8}$ or $6\frac{1}{8}$ or Hence the turning point of f is $\left(-\frac{5}{4}; \frac{49}{8}\right)$ or $(-1,25; 6,13)$ OR $f(x) = -2x^2 - 5x + 3$ $= -2\left(x^2 + \frac{5x}{2} - \frac{3}{2}\right)$ $= -2\left(x + \frac{5}{4}\right)^2 + \frac{25}{8} + \frac{24}{8}$ $= -2\left(x + \frac{5}{4}\right)^2 + \frac{49}{8}$ Hence the turning point of f is $\left(-\frac{5}{4}; \frac{49}{8}\right)$	$f'(x) = 0$ $-4x - 5 = 0$ $x = -\frac{5}{4}$ or $-1\frac{1}{4}$	✓ $x = -\frac{(-5)}{2(-2)}$ or $\frac{1}{2} - 3$ $f'(x) = 0$ or $\frac{1}{2} - 3$ ✓ x -coordinate ✓ y -coordinate (3) ✓ $-2\left(x + \frac{5}{4}\right)^2 + \frac{49}{8}$ ✓ x -coordinate ✓ y -coordinate (3)

4.4		✓ shape ✓ intercepts with the axes ✓ turning point (3) [10]
-----	--	---


QUESTION/VRAAG 5

Given: $g(x) = k^x$ and $y = g^{-1}(x)$ 		
5.1.1	$k^2 = 36$ $k = 6$	✓ $k^2 = 36$ ✓ answer (2)
5.1.2	g: $y = 6^x$ g^{-1}: $x = 6^y$ $y = \log_6 x$ or $y = \frac{\log x}{\log 6}$ NOTE: Answer only : 2/2 marks	✓ $x = 6^y$ ✓ $y = \log_6 x$ or $y = \frac{\log x}{\log 6}$ (2)
5.1.3	$0 < x \leq 1$ OR $(0; 1]$	✓ $0 < x$ ✓ $x \leq 1$ (2)
5.1.4	$x > 3$ OR $(3; \infty)$	✓ answer (1)

5.2.1		✓ vertical line ✓ vertical line has x-intercept $(1 ; 0)$ (2) NOTE: Any other graph cutting at $(1 ; 0)$: 0 / 2 marks
5.2.2	NOTE: If $x = 1$ is not drawn in 5.2.1: 0/2 marks No, the inverse of f is not a function because the vertical line test fails in the graph of $x = 1$. <i>Nee, die inverse van f is nie 'n funksie nie omdat die vertikalelyn-toets vir die grafiek $x = 1$ faal.</i> OR No, the inverse of f is not a function because the horizontal line test fails in the graph of $y = 1$. <i>Nee, die inverse van f is nie 'n funksie nie omdat die horisontalelyn-toets vir die grafiek $y = 1$ faal.</i> OR Since f as defined is a many-to-one function it does not have an inverse function. <i>f is gedefinieerd as 'n meer-tot-een-funksie en het dus nie 'n inverse funksie nie.</i> OR The inverse of f is a one-to-many relation. <i>Die inverse funksie van f is 'n een-tot-meer relasie.</i> OR	✓ not a function ✓ valid reason (2) ✓ not a function ✓ valid reason (2) ✓ not a function ✓ valid reason (2) ✓ not a function ✓ explanation (2)

	No, the inverse of f is not a function because there are some input (x) values (for example, $x = 0$) which have more than one output (y) value. <i>Nee, die inverse van f is nie 'n funksie nie omdat van die x-waardes (bv. $x = 0$) meer as een y-waarde het.</i> OR No, for one x-value there are more than one y-values. <i>Nee, vir 'n x-waarde is daar meer as een y-waarde.</i>	✓ not a function ✓ valid reason (2) ✓ not a function ✓ valid reason (2) [11]
--	---	--

QUESTION/VRAAG 6

	Given: $f(x) = \frac{x-d}{x-p}$ 	
6.1.1	$0 = \frac{2-d}{2-p}$ $d = 2$ $-1-p = 0$ $p = -1$ NOTE: If candidate leaves answer as $f(x) = \frac{x-2}{x+1}$: 2/2 marks	✓ d value ✓ p value (2)
6.1.2	$y = \frac{x-2}{x+1}$ $= \frac{(x+1)-3}{x+1}$ $= \frac{x+1}{x+1} - \frac{3}{x+1}$ $= \frac{-3}{x+1} + 1$ OR $\frac{-3}{x+1} + 1$ $= \frac{-3+x+1}{x+1}$ $= \frac{x-2}{x+1}$ NOTE: - If the candidate starts with $y = \frac{a}{x+1} + 1$ and substitutes (2 ; 0) and proves $a = -3$: 0/2 marks - If the candidate starts with $y = \frac{-3}{x+1} + 1$ and calculates (2 ; 0) as x-intercept: 0/2 marks	✓ $\frac{x-2}{x+1} = \frac{x+1-3}{x+1}$ ✓ $\frac{x+1}{x+1} - \frac{3}{x+1}$ (2)

	OR $\begin{array}{r} 1 \\ x+1 \overline{)x-2} \\ \underline{-x-1} \\ -3 \end{array}$ Remainder = -3 $\therefore x-2 = 1(x+1) - 3$ $\frac{x-2}{x+1} = 1 - \frac{3}{x+1}$ $f(x) = \frac{-3}{x+1} + 1$	✓ $\frac{-3+x+1}{x+1}$ ✓ simplification (2) ✓ long division ✓ remainder = -3 (2)
6.1.3	P(-1 ; 1)	✓ x-coordinate ✓ y-coordinate (2)
6.1.4	(-2 ; 4)	✓ x-coordinate ✓ y-coordinate (2)
6.2	$q = 1$ Substitute A(0; -2) into $g(x) = p \cdot 2^x + 1$: $-2 = p(2)^0 + 1$ $p = -3$ Hence $g(x) = -3 \cdot 2^x + 1$ NOTE: Answer only: 3/3 marks	✓ $q = 1$ ✓ substitute A(0; -2) ✓ $p = -3$ (3) [11]

QUESTION/VRAAG 7

7.1.1	$1 + i_{eff} = \left(1 + \frac{0,09}{4}\right)^4$ $i_{eff} = \left(1 + \frac{0,09}{4}\right)^4 - 1$ $= 0,093083318$ $= 0,0931$ $= 9,31\%$	$\checkmark \left(1 + \frac{0,09}{4}\right)^4$ $\checkmark 9,31\% \text{ or } 0,0931$ (2)
7.1.2	$A = P(1+i)^n$ $30\,440 = 12\,500\left(1 + \frac{0,09}{4}\right)^{4k}$ $\frac{30440}{12500} = \left(1 + \frac{0,09}{4}\right)^{4k}$ $2,4352 = 1,0225^{4k}$ $4k = \log_{1,0225} 2,4352$ $4k = 40,00020365\dots$ $k = 10 \text{ years}$ $A = P(1+i)^n$ $30\,440 = 12\,500\left(1 + \frac{0,09}{4}\right)^{4k}$ $\frac{30440}{12500} = \left(1 + \frac{0,09}{4}\right)^{4k}$ $2,4352 = 1,0225^{4k}$ $4k \log 1,0225 = \log 2,4352$ $4k = \frac{\log 2,4352}{1,0225}$ $4k = 40,00020365\dots$ $k = 10 \text{ years}$ OR $A = P(1+i)^n$ $30\,440 = 12\,500(1 + 0,09308\dots)^k$ $\frac{30440}{12500} = (1 + 0,09308\dots)^k$ $2,4352 = (1,09308\dots)^k$ $k = \log_{1,09308\dots} 2,4352$ $k = 9,998336572\dots$ $k = 10 \text{ years}$ $A = P(1+i)^n$ $30\,440 = 12\,500(1 + 0,09308\dots)^k$ $\frac{30440}{12500} = (1 + 0,09308\dots)^k$ $2,4352 = (1,09308\dots)^k$ $k \log 1,09308\dots = \log 2,4352$ $k = \frac{\log 2,4352}{\log 1,09308\dots}$ $k = 9,998336572\dots$ $k = 10 \text{ years}$ OR NOTE: Incorrect formula: max 2/5 marks If A and P are swapped: max 2 / 5 marks	$\checkmark n = 4k$ $\checkmark i = \frac{0,09}{4}$ $\checkmark \text{subs into correct formula}$ $\checkmark \text{use of logs}$ $\checkmark \text{answer}$ (5) $\checkmark n = k$ $\checkmark i = 0,09308\dots \text{ (from 7.1.1)}$ $\checkmark \text{subs into correct formula}$ $\checkmark \text{use of logs}$ $\checkmark \text{answer}$ (5)


	$4n = \frac{\log \frac{A}{P}}{\log(1+i)}$ $4k = \frac{\log \frac{30440}{12500}}{\log\left(1 + \frac{0,09}{4}\right)}$ $4k = 40,00020365\dots$ $k = 10 \text{ years}$	✓ $n = 4k$ ✓ $i = \frac{0,09}{4}$ ✓ subs into correct formula ✓ use of logs ✓ answer (5)
7.2.1	30% of R18 480 $= \left(\frac{30}{100}\right)(18\,480)$ $= \text{R } 5\,544$	✓ answer (1)
7.2.2	$P = \frac{x[1 - (1+i)^{-n}]}{i}$ $= \frac{5544 \left[1 - \left(1 + \frac{0,08}{12}\right)^{-300} \right]}{\frac{0,08}{12}}$ $= \text{R } 718\,305,71$ OR $x \left(1 + \frac{0,08}{12}\right)^{300} = \frac{5544 \left[\left(1 + \frac{0,08}{12}\right)^{300} - 1 \right]}{\frac{0,08}{12}}$ $x \left(1 + \frac{0,08}{12}\right)^{300} = 5272490,33\dots$ $x = \text{R } 718\,305,71$ NOTE: Incorrect formula: max 2/4 marks If the answer is R 717 550,42 (used $n = 299$): award 3 /4 marks	✓ $i = \frac{0,08}{12}$ ✓ $n = 300$ ✓ substitution ✓ answer (4) ✓ $i = \frac{0,08}{12}$ ✓ $n = 300$ ✓ substitution ✓ answer (4) [12]

QUESTION/VRAAG 8

8.1.1	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{3(x+h)^2 - 4 - (3x^2 - 4)}{h}$ $= \lim_{h \rightarrow 0} \frac{3x^2 + 6xh + 3h^2 - 4 - 3x^2 + 4}{h}$ $= \lim_{h \rightarrow 0} \frac{6xh + 3h^2}{h}$ $= \lim_{h \rightarrow 0} \frac{h(6x + 3h)}{h} ; h \neq 0$ $= \lim_{h \rightarrow 0} (6x + 3h)$ $= 6x$ OR $f(x) = 3x^2 - 4$ $f(x+h) = 3(x+h)^2 - 4$ $= 3x^2 + 6xh + 3h^2 - 4$ $f(x+h) - f(x) = 6xh + 3h^2$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{6xh + 3h^2}{h}$ $= \lim_{h \rightarrow 0} \frac{h(6x + 3h)}{h}$ $= \lim_{h \rightarrow 0} (6x + 3h)$ $= 6x$ NOTE: - Incorrect notation: max 4/5 marks (Leaving out limit / incorrect use of limit, leaving out $f'(x)$, = in the wrong place constitute incorrect notation) - If $\lim_{h \rightarrow 0} \frac{3x^2 + 6xh + 3h^2 - 3x^2 - 8}{h}$, take out common factor, then correct to the final answer: max 3/5 marks. NOTE: If candidate uses differentiation rules: 0/5 marks	✓ formula ✓ substitution of $x + h$ ✓ simplification to $\frac{6xh + 3h^2}{h}$ ✓ $\lim_{h \rightarrow 0} (6x + 3h)$ ✓ answer (5) ✓ substitution of $x + h$ ✓ simplification to $6xh + 3h^2$ ✓ formula ✓ $\lim_{h \rightarrow 0} (6x + 3h)$ ✓ answer (5)
8.1.2	$f(x) = 3x^2 - 4$ average gradient of f between $A(-2; y)$ and $B(x; 23)$ $y = 3(-2)^2 - 4 = 8$ $23 = 3x^2 - 4$ $27 = 3x^2$ $9 = x^2$ $x = 3$	✓ $y = 8$ ✓ $23 = 3x^2 - 4$ ✓ $x = 3$


	Average gradient $= \frac{23 - y}{x - (-2)}$ $= \frac{23 - 8}{3 + 2}$ $= 3$	✓ $\frac{23 - y}{x - (-2)}$ ✓ answer (5)
8.2	$y = \frac{x + 5}{x^{\frac{1}{2}}}$ $= \frac{x}{x^{\frac{1}{2}}} + \frac{5}{x^{\frac{1}{2}}}$ $= x^{\frac{1}{2}} + 5x^{-\frac{1}{2}}$ $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} - \frac{5}{2}x^{-\frac{3}{2}}$ OR $y = \frac{x + 5}{x^{\frac{1}{2}}}$ By the quotient rule $\frac{dy}{dx} = \frac{1 \cdot x^{\frac{1}{2}} - \frac{1}{2}x^{-\frac{1}{2}}(x + 5)}{(x^{\frac{1}{2}})^2}$ $= \frac{x^{\frac{1}{2}} - \frac{1}{2}x^{\frac{1}{2}} - \frac{5}{2}x^{-\frac{1}{2}}}{x}$ $= \frac{1}{2x^{\frac{1}{2}}} - \frac{5}{2x^{\frac{3}{2}}}$	NOTE: There is a maximum penalty of 1 mark for incorrect notation in question 8. ✓ $x^{\frac{1}{2}} + 5x^{-\frac{1}{2}}$ ✓ $\frac{1}{2}x^{-\frac{1}{2}}$ or $\frac{1}{2\sqrt{x}}$ ✓ $-\frac{5}{2}x^{-\frac{3}{2}}$ or $\frac{-5}{2\sqrt{x^3}}$ (3) ✓ ✓ $\frac{1 \cdot x^{\frac{1}{2}} - \frac{1}{2}x^{-\frac{1}{2}}(x + 5)}{(x^{\frac{1}{2}})^2}$ ✓ $\frac{1}{2}x^{-\frac{1}{2}}$ or $\frac{1}{2\sqrt{x}}$ or $-\frac{5}{2}x^{-\frac{3}{2}}$ or $\frac{-5}{2\sqrt{x^3}}$ (3)
8.3	$f(x) = -3x^3 - 4x + 5$ $f'(x) = -9x^2 - 4$ $m_{\tan} = -9(-1)^2 - 4$ $= -13$	✓ $-9x^2$ ✓ -4 ✓ substitution of $x = -1$ ✓ answer (4) [17]

QUESTION/VRAAG 9

	Given: $f(x) = x^3 + ax^2 + bx - 2$ $P(-1; -1)$ and R are the turning points of f. 	
9.1	$f(x) = x^3 + ax^2 + bx - 2$ $-1 = (-1)^3 + a(-1)^2 + b(-1) - 2$ $2 = a - b \quad \dots(1)$ $f'(x) = 3x^2 + 2ax + b$ $0 = 3(-1)^2 + 2a(-1) + b$ $-3 = -2a + b \quad \dots(2)$ $-1 = -a \quad \dots(1) + (2)$ $a = 1$ $b = -1$ NOTE: - The $= 0$ must be explicitly stated - If the candidate starts with what is given and then prove $(1; 1)$ is a TP: 0/6 marks	$\checkmark -1 = (-1)^3 + a(-1)^2 + b(-1) - 2$ $\checkmark 2 = a - b$ $\checkmark f'(x) = 3x^2 + 2ax + b$ $\checkmark f'(-1) = 0$ $\checkmark -3 = -2a + b$ $\checkmark$ method (6)
9.2	R is a turning point of f, hence at R, $f'(x) = 0$ i.e. $3x^2 + 2x - 1 = 0$ $(3x - 1)(x + 1) = 0$ NOTE: Answer only: 1/3 marks $x = \frac{1}{3} \text{ or } -1$ $\therefore x = \frac{1}{3}$	$\checkmark f'(x) = 0$ $\checkmark f'(x) = 3x^2 + 2x - 1$ $\checkmark$ selection of $x = \frac{1}{3}$ (3)
9.3	$(-1; 2f(-1) - 4)$ $= (-1; -6)$ OR $\left(\frac{1}{3}; 2f\left(\frac{1}{3}\right) - 4\right)$ $= \left(\frac{1}{3}; -\frac{226}{27}\right) \text{ or } (0,33; -8,37)$	$\checkmark$ x-coordinate $\checkmark$ y-coordinate (2) $\checkmark$ x-coordinate $\checkmark$ y-coordinate (2) [11]

10.1	$\frac{dr}{dt} = -0,4t + 10$ $0 = -0,4t + 10$ $0,4t = 10$ $t = \frac{10}{0,4}$ $= 25 \text{ seconds}$ OR $t = -\frac{b}{2a}$ $= -\frac{10}{2(-0,2)}$ $= 25 \text{ seconds}$ OR $r = \frac{1}{5}t(50 - t)$ $0 = \frac{t}{5}(50 - t)$ $t = 0 \text{ or } t = 50$ Fastest at $t = \frac{0 + 50}{2}$ $t = 25 \text{ seconds}$	$\checkmark \frac{dr}{dt} = -0,4t + 10$ $\checkmark 0 = -0,4t + 10$ $\checkmark t \text{ value}$ $\checkmark t = -\frac{b}{2a}$ $\checkmark \text{ substitution}$ $\checkmark t \text{ value}$ $\checkmark t = 0 \text{ or } t = 50$ $\checkmark t = \frac{0 + 50}{2}$ $\checkmark t \text{ value}$
10.2	$-0,2t^2 + 10t = 0$ $t(-0,2t + 10) = 0$ $-0,2t + 10 = 0 \quad \text{or} \quad t = 0$ $t = \frac{-10}{-0,2}$ $= 50 \text{ sec}$ Hence the water stops flowing 50 seconds after it started. OR $-0,2t^2 + 10t = 0$ $t^2 - 50t = 0$ $t(t - 50) = 0$ $t = 0 \text{ or } t = 50$ Hence the water stops flowing 50 seconds after it started. NOTE: Answer only: 3/3 marks	$\checkmark -0,2t^2 + 10t = 0$ $\checkmark \text{ factors}$ $\checkmark \text{ answer}$ $\checkmark -0,2t^2 + 10t = 0$ $\checkmark \text{ factors}$ $\checkmark \text{ answer}$

QUESTION/VRAAG 11

11.1	$y \geq 50$ $x \leq 80$ $y \leq 5x$ OR $\frac{y}{5} \leq x$ $y \leq -\frac{5}{4}x + 200$ OR $5x + 4y \leq 800$	$\checkmark y \geq 50$ $\checkmark x \leq 80$ $\checkmark y \leq 5x$ $\checkmark 5x + 4y \leq 800$ (4)
11.2		$\checkmark y = 50$ $\checkmark x = 80$ $\checkmark 5x + 4y = 800$ $\checkmark y = 5x$ $\checkmark$ feasible region / gangbare gebied (5) NOTE: Maximum of 2 marks for vertical and horizontal lines
11.3.1	$P = 30x + 20y$	$\checkmark P = 30x + 20y$ (1)
11.3.2	$y = -\frac{3}{2}x + \frac{P}{20}$ 80 short-sleeved shirts and 100 long-sleeved shirts (Point (80 ; 100))	$\checkmark$ 80 short-sleeved shirts $\checkmark$ 100 long-sleeved shirts (2)
11.4	$P = ax + by$ $m = -\frac{a}{b}$ $-\frac{5}{4} = -\frac{a}{b}$ $\frac{a}{b} = \frac{5}{4}$	NOTE: Answer only: 2/2 marks $\checkmark -\frac{5}{4} = -\frac{a}{b}$ $\checkmark$ answer (2) [14]

TOTAL/TOTAAL: 150

