




eduresourceza.com



Education and Sport Development

Department of Education and Sport Development
Departement van Onderwys en Sportontwikkeling
Lefapha la Thuto le Tlhabololo ya Metshameko

NORTH WEST PROVINCE

NATIONAL SENIOR CERTIFICATE

GRADE 12

MATHEMATICS P1

SEPTEMBER 2018

MEMORANDUM

MARKS: 150

This memorandum consists of 17 pages.

NOTE:

- If a candidate answered a question TWICE, only mark the FIRST attempt.
- Consistent accuracy applies in ALL aspects of the marking memorandum.

QUESTION 1

1.1.1	$2x(5x - 3) = 0$ $2x = 0$ or $5x - 3 = 0$ $x = 0$ $5x = 3$ $x = \frac{3}{5}$	$\checkmark x = 0$ $\checkmark x = \frac{3}{5}$ (2)
1.1.2	$-x^2 + 4 = 5x$ $-x^2 - 5x + 4 = 0$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{5 \pm \sqrt{(-5)^2 - 4(-1)(4)}}{2(-1)}$ $= \frac{5 \pm \sqrt{41}}{-2}$ $x = -5,70$ or $x = 0,70$	$\checkmark$ standard form $\checkmark$ substitution into the correct formula $\checkmark x = -5,70$ $\checkmark x = 0,70$ (4)
1.1.3	$\sqrt{x-6} - 2 = \frac{15}{\sqrt{x-6}}$ Let $k = \sqrt{x-6}$ $k - 2 = \frac{15}{k}$ $k^2 - 2k - 15 = 0$ $(k-5)(k+3) = 0$ $k = 5$ or $k = -3$ $\sqrt{x-6} = 5$ or $\sqrt{x-6} = -3$ $x - 6 = 25$ $n.a.$ $x = 31$	$\checkmark k^2 - 2k - 15 = 0$ $\checkmark (k-5)(k+3) = 0$ $\checkmark k = 5$ or $k = -3$ $\checkmark \sqrt{x-6} = -3$ n.a. $\checkmark x = 31$ (5)
1.1.4	$(x^2 + 2)(x - 3) < 0$ $(x^2 + 2) > 0$ $\therefore (x - 3) < 0$ $x < 3$	$\checkmark (x - 3) < 0$ $\checkmark x < 3$ (2)

1.2	$x + 2y = 3$ $x = 3 - 2y \dots\dots\dots(1)$ $3x^2 + 4xy + 9y^2 - 16 = 0 \dots\dots\dots(2)$ $3(3 - 2y)^2 + 4(3 - 2y)y + 9y^2 - 16 = 0$ $3(9 - 12y + 4y^2) + 12y - 8y^2 + 9y^2 - 16 = 0$ $27 - 36y + 12y^2 + 12y - 8y^2 + 9y^2 - 16 = 0$ $13y^2 - 24y + 11 = 0$ $(13y - 11)(y - 1) = 0$ $y = \frac{11}{13} \quad \text{or} \quad y = 1$ $x = 3 - 2\left(\frac{11}{13}\right) \quad \text{or} \quad x = 3 - 2(1)$ $= \frac{17}{13} \quad \quad \quad = 1$ OR $x + 2y = 3$ $2y = 3 - x$ $y = \frac{3 - x}{2} \dots\dots\dots(1)$ $3x^2 + 4xy + 9y^2 - 16 = 0 \dots\dots\dots(2)$ $3x^2 + 4x\left(\frac{3 - x}{2}\right) + 9\left(\frac{3 - x}{2}\right)^2 - 16 = 0$ $3x^2 + 6x - 2x^2 + \frac{9(9 - 6x + x^2)}{4} - 16 = 0$ $12x^2 + 24x - 8x^2 + 81 - 54x + 9x^2 - 64 = 0$ $13x^2 - 30x + 17 = 0$ $(13x - 17)(x - 1) = 0$ $x = \frac{17}{13} \quad \text{or} \quad x = 1$ $y = \frac{3 - \frac{17}{13}}{2} \quad \text{or} \quad y = \frac{3 - 1}{2}$ $= \frac{11}{13} \quad \quad \quad = 1$	$\checkmark x = 3 - 2y$ $\checkmark \text{ substitution}$ $\checkmark \text{ standard form}$ $\checkmark \text{ factors or formula}$ $y = \frac{24 \pm \sqrt{(-24)^2 - 4(13)(11)}}{2(13)}$ $\checkmark \text{ both } y\text{-values}$ $\checkmark \text{ both } x\text{-values}$ (6) $\checkmark y = \frac{3 - x}{2}$ $\checkmark \text{ substitution}$ $\checkmark \text{ standard form}$ $\checkmark \text{ factors or formula}$ $x = \frac{30 \pm \sqrt{(-30)^2 - 4(13)(17)}}{2(13)}$ $\checkmark \text{ both } x\text{-values}$ $\checkmark \text{ both } y\text{-values}$ (6)
-----	--	--

1.3	$2^{2022} \cdot 5^{2018}$ $= 2^{2018+4} \cdot 5^{2018}$ $= 2^{2018} \cdot 2^4 \cdot 5^{2018}$ $= (2.5)^{2018} \cdot 16$ $= 10^{2018} \cdot 16$ $= (100000000.....)(16)$ $= (160000000.....)$ Sum of the digits = $1 + 6 = 7$	✓ $2^{2018} \cdot 2^4 \cdot 5^{2018}$ ✓ $(2.5)^{2018} \cdot 16$ ✓ $(160000000.....)$ ✓ Sum of the digits = 7 (4) [23]
-----	--	---


QUESTION 2

2.1.1	$\frac{T_2}{T_1} = \frac{T_3}{T_2}$ $\frac{x-1}{x+1} = \frac{2x-5}{x-1}$ $(x-1)^2 = (2x-5)(x+1)$ $x^2 - 2x + 1 = 2x^2 - 3x - 5$ $0 = x^2 - x - 6$ $0 = (x-3)(x+2)$ $x = 3 \text{ or } x = -2$	✓ $\frac{x-1}{x+1} = \frac{2x-5}{x-1}$ ✓ simplification ✓ standard form ✓ factors ✓ both answers (5)
2.1.2	If $x = 3$: If $x = -2$: $R_1: 4; 2; 1$ $R_2: -1; -3; -9$ $r = \frac{1}{2}$ $r = 3$ For convergent: $-1 < r < 1$ $\therefore x = 3$	✓✓ both r-values ✓ $-1 < r < 1$ ✓ $x = 3$ (4)
2.1.3	$S_{\infty} = \frac{a}{1-r}$ $= \frac{4}{1-\frac{1}{2}}$ $= 8$	✓ substitution ✓ answer (2)

2.2	$1 + 2 + 4 + \dots$ (30 terms) $S_n = \frac{a(r^n - 1)}{r - 1}$ $S_{30} = \frac{1(2^{30} - 1)}{2 - 1}$ $= 1\,073\,741\,823$ $\therefore$ the 2 nd option is the best	✓ substitution ✓ answer ✓ option 2 (3) [14]
-----	---	--


QUESTION 3

3.1	
-----	--

	Sequence of last chair of each row:  $2a = 2$ $a = 1$ $T_2 - T_1 = 3a + b$ $7 = 3(1) + b$ $b = 4$ $T_1 = a + b + c$ $5 = 1 + 4 + c$ $c = 0$ $T_n = n^2 + 4n$ $T_{25} = (25)^2 + 4(25)$ $= 725$ $T_{24} = (24)^2 + 4(24)$ $= 672$ $\therefore \text{Number of chairs in row } 25 = 725 - 672$ $= 53$	✓ general term ✓ answer of both terms ✓ answer (3)
3.4	Sequence of number of chairs per row : 5; 7; 9; 11 $S_n = \frac{n}{2}[2a + (n-1)d]$ $2\,000 = \frac{n}{2}[2(5) + (n-1)(2)]$ $2\,000 = 5n + n^2 - n$ $0 = n^2 + 4n - 2\,000$ $n = \frac{-4 \pm \sqrt{4^2 - 4(1)(-2\,000)}}{2(1)}$ $n = -46,77 \quad \text{or} \quad n = 42,77$ $n.a. \quad n = 42$ $\therefore$ 42 complete rows OR	✓ substitution ✓ standard form ✓ substitution into formula ✓ $n = 42,77$ ✓ $n = 42$ (5)

Last chair of each row : 5; 7; 9; 11 $T_n = n^2 + 4n$ $2000 = n^2 + 4n$ $0 = n^2 + 4n - 2000$ $n = \frac{-4 \pm \sqrt{4^2 - 4(1)(-2000)}}{2(1)}$ $n = -46,77 \quad \text{or} \quad n = 42,77$ $n.a. \quad n = 42$ $\therefore 42 \text{ complete rows}$	✓ substitution ✓ standard form ✓ substitution into formula ✓ $n = 42,77$ ✓ $n = 42$ (5) [14]
--	---


QUESTION 4

4.1	$f(x) = \frac{a}{x-2} + 3$ $0 = \frac{a}{-1-2} + 3$ $-3 = \frac{a}{-3}$ $9 = a$ $f(x) = \frac{9}{x-2} + 3$	✓ $f(x) = \frac{a}{x-2} + 3$ ✓ Substitution B(-1; 0) ✓ $a = 9$ (3)
4.2	$y = \frac{9}{0-2} + 3$ $= \frac{9}{-2} + 3$ $= -\frac{3}{2} \quad \therefore \left(0; -\frac{3}{2}\right) \quad \text{OR} \quad y = -1\frac{1}{2} \quad \therefore \left(0; -1\frac{1}{2}\right)$	✓ $x = 0$ ✓ answer (2)
4.3		✓ x- and y-intercepts ✓ both asymptotes ✓ form (3) [8]

QUESTION 5

5.1	$x = -1$	✓ $x = -1$ (1)
5.2	$y = \frac{2}{0+1} - 3$ $= 2 - 3$ $= -1$ $\therefore P(0; -1)$	✓ $x = 0$ ✓ $y = -1$ (2)
5.3	$f(x) = a(x+1)^2 - 3$ $-1 = a(0-1)^2 - 3$ $a = 2$ $f(x) = 2(x+1)^2 - 3$	✓ $f(x) = a(x+1)^2 - 3$ ✓ $-1 = a(0-1)^2 - 3$ ✓ $a = 2$ (3)
5.4	$h(x) = -(x+1) - 3$ $= -x - 1 - 3$ $= -x - 4$ OR $h(x) = -x + c$ $-3 = -(-1) + c$ $c = -4$ $h(x) = -x - 4$	✓ $h(x) = -(x+1) - 3$ ✓ answer (2) ✓ $-3 = -(-1) + c$ ✓ answer (2)
5.5	$h(x) = -x - 4$ $k > 3$	✓✓ $k > 3$ (2)
5.6	$m(x) = \frac{2}{2x+1} - 3 + 5$ $= \frac{2}{2\left(x + \frac{1}{2}\right)} + 2$ $= \frac{1}{x + \frac{1}{2}} + 2$ $x \in \mathbb{R}; x \neq -\frac{1}{2}$	✓ $m(x) = \frac{2}{2x+1} - 3 + 5$ ✓ $\frac{1}{x + \frac{1}{2}} + 2$ ✓ $x \in \mathbb{R}; x \neq -\frac{1}{2}$ (3) [13]

QUESTION 6

6.1	$g(x) = a^x$ $3 = a^{-1}$ $3 = \frac{1}{a}$ $a = \frac{1}{3}$	$\checkmark 3 = a^{-1}$ (1)
6.2	$[-2; 0)$ OR $-2 \leq x < 0$	$\checkmark\checkmark [-2; 0)$ OR $\checkmark\checkmark -2 \leq x < 0$ (2)
6.3	$f: y = \frac{1}{4}x^2; x \geq -2; y \geq 0$ $f^{-1}: x = \frac{1}{4}y^2$ $4x = y^2$ $y = \pm \sqrt{4x} = \pm 2\sqrt{x}; x \geq 0$	$\checkmark$ swop x and y $\checkmark y = \pm \sqrt{4x} = \pm 2\sqrt{x}$ (2)
6.4		$\checkmark$ form $\checkmark$ y -value of end point (2)
6.5	$y \geq -2$	$\checkmark$ notation $\checkmark -2$ (2)
6.6	$g: y = \left(\frac{1}{3}\right)^x$ $g^{-1}: x = \left(\frac{1}{3}\right)^y$ $y = \log_{\frac{1}{3}} x; x > 0$	$\checkmark$ swop x and y $\checkmark y = \log_{\frac{1}{3}} x$ (2)
6.7	$x \in (0; 3]$	$\checkmark (0;$ $\checkmark 3]$ (2) [13]

QUESTION 7

7.1	$P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $2\,000\,000 = \frac{35\,000 \left[1 - \left(1 + \frac{0,093}{12} \right)^{-n} \right]}{\frac{0,093}{12}}$ $\frac{31}{70} = 1 - \left(1 + \frac{0,093}{12} \right)^{-n}$ $\left(1 + \frac{0,093}{12} \right)^{-n} = \frac{39}{70} \quad \text{OR} \quad \log \left(1 + \frac{0,093}{12} \right)^{-n} = \log \frac{39}{70}$ $-n = \log_{\left(1 + \frac{0,093}{12} \right)} \frac{39}{70} \quad -n = \frac{\log \frac{39}{70}}{\log \left(1 + \frac{0,093}{12} \right)}$ $n = 75,77$ $\therefore$ He will be able to live for 75 months from his investment	✓ substitution of i ✓ substitution into correct formula ✓ simplification ✓ correct use of log ✓ answer (5)
7.2.1	$A = P(1 + i)^n$ $= 1\,200\,000(1 + 0,075)^6$ $= R1\,851\,961,83$ He will need = $R1\,851\,961,83 - 400\,000$ $= R1\,451\,961,83$	✓ substitution ✓ R1 851 961,83 ✓ answer (3)
7.2.2	$A = P(1 + i)^n$ $1\,451\,961,83 = P \left(1 + \frac{0,11}{4} \right)^2$ $P = 1\,375\,281,30$ $F = \frac{x[(1 + i)^n - 1]}{i}$ $1\,375\,281,30 = \frac{x \left[\left(1 + \frac{0,11}{4} \right)^{21} - 1 \right]}{\frac{0,11}{4}}$ $x = R49\,261,76$ OR	✓ substitution ✓ substitution of i ✓ substitution of n ✓ substitution into correct formula ✓ answer (5)

	$F = \frac{x(1+i)^n[(1+i)^n - 1]}{i}$ $1\,451\,961,83 = \frac{x\left(1 + \frac{0,11}{4}\right)^2 \cdot \left[\left(1 + \frac{0,11}{4}\right)^{21} - 1\right]}{\frac{0,11}{4}}$ $x = R\,49\,261,76$	✓ $\left(1 + \frac{0,11}{4}\right)^2$ ✓ substitution of i ✓ substitution of n ✓ substitution into correct formula ✓ answer
7.2.3	Quarterly deposit to withdraw R 8 000 at the end of each year from year 2 to year 5: $F = \frac{x[(1+i)^n - 1]}{i}$ $8\,000 = \frac{x\left[\left(1 + \frac{0,11}{4}\right)^4 - 1\right]}{\frac{0,11}{4}}$ $x = R\,1\,919,36$ New quarterly deposit $= 49\,261,76 + 1\,919,36$ $= R\,51\,181,12$	✓ substitution of n ✓ substitution into correct formula ✓ $x = R\,1\,919,36$ ✓ answer

(5)

(4)
[17]

QUESTION 8


8.1	$f(x) = 2x^2 - 5x + 3$ $f(x+h) = 2(x+h)^2 - 5(x+h) + 3$ $= 2(x^2 + 2xh + h^2) - 5x - 5h + 3$ $= 2x^2 + 4xh + 2h^2 - 5x - 5h + 3$ $f(x+h) - f(x) = 2x^2 + 4xh + 2h^2 - 5x - 5h + 3$ $\quad - (2x^2 - 5x + 3)$ $= 4xh + 2h^2 - 5h$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - 5h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 5)}{h}$ $= \lim_{h \rightarrow 0} (4x + 2h - 5)$ $= 4x - 5$	✓ $2(x+h)^2 - 5(x+h) + 3$ ✓ $4xh + 2h^2 - 5h$ ✓ formula ✓ factor ✓ answer
-----	--	---

(5)

	OR $f(x) = 2x^2 - 5x + 3$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{2(x+h)^2 - 5(x+h) + 3 - (2x^2 - 5x + 3)}{h}$ $= \lim_{h \rightarrow 0} \frac{2(x^2 + 2xh + h^2) - 5x - 5h + 3 - 2x^2 + 5x - 3}{h}$ $= \lim_{h \rightarrow 0} \frac{2x^2 + 4xh + 2h^2 - 5x - 5h + 3 - 2x^2 + 5x - 3}{h}$ $= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - 5h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 5)}{h}$ $= \lim_{h \rightarrow 0} (4x + 2h - 5)$ $= 4x - 5$	✓ formula ✓ $2(x+h)^2 - 5(x+h) + 3$ ✓ $4xh + 2h^2 - 5h$ ✓ factor ✓ answer (5)
8.2	$y = \frac{2x^2}{3\sqrt{x}} - \frac{2x^3 + 1}{x^3}$ $= \frac{2x^2}{3x^{\frac{1}{2}}} - 2 - \frac{1}{x^3}$ $= \frac{2}{3}x^{\frac{3}{2}} - 2 - x^{-3}$ $\frac{dy}{dx} = x^{\frac{1}{2}} + 3x^{-4}$	✓ $\frac{2}{3}x^{\frac{3}{2}}$ ✓ -2 ✓ $-x^{-3}$ ✓ $x^{\frac{1}{2}}$ ✓ $3x^{-4}$ (5) [10]

QUESTION 9


9.1.1	$f(x) = -2x^3 + 5x^2 + 4x - 3$ $0 = (x-3)(-2x^2 - x + 1)$ $x-3=0 \quad \text{or} \quad -2x^2 - x + 1 = 0$ $x=3 \quad \quad \quad 2x^2 + x - 1 = 0$ $(3; 0) \quad \quad (2x-1)(x+1) = 0$ $2x=1 \quad \text{or} \quad x=-1$ $x = \frac{1}{2} \quad \quad (-1; 0)$ $\left(\frac{1}{2}; 0\right)$	✓ $-2x^2 - x + 1$ ✓ (3; 0) ✓ factors or formula ✓ both coordinates $\left(\frac{1}{2}; 0\right) \quad (-1; 0)$ (4)
-------	---	--

9.1.2	$f'(x) = -6x^2 + 10x + 4$ $0 = -6x^2 + 10x + 4$ $3x^2 - 5x - 2 = 0$ $(3x + 1)(x - 2) = 0$ or $x = \frac{-10 \pm \sqrt{10^2 - 4(-6)(4)}}{2(-6)}$ $3x = -1$ or $x = 2$ $x = -\frac{1}{3}$	✓ $-6x^2 + 10x + 4$ ✓ $f'(x) = 0$ ✓ factors or formula ✓ both x answers (4)
9.1.3	$f''(x) = -12x + 10$ $0 = -12x + 10$ $12x = 10$ $x = \frac{10}{12} = \frac{5}{6}$ $\therefore x < \frac{5}{6}$	✓ $0 = -12x + 10$ ✓ $x < \frac{5}{6}$ (2)
9.2.1		✓ form ✓ both x -intercepts ✓ x -value of both turning points ✓ x -value of point of inflection (4)
9.2.2	$x < -2$ or $x > 2$; $x \neq 4$	✓ $x < -2$ ✓ $x > 2$ ✓ $x \neq 4$ (3) [17]

QUESTION 10

10.1	$N(t) = t^3 - 12t^2 + 36t + 8$ $N(0) = 8$ $\therefore 8$ people	✓ answer (1)
10.2	$N'(t) = 3t^2 - 24t + 36$ increasing $N'(t) \geq 0$ $3t^2 - 24t + 36 \geq 0$ $t^2 - 8t + 12 \geq 0$ $(t - 6)(t - 2) \geq 0$ $t \geq 6$ or $t \leq 2$ $\therefore$ for first 2 hours after opening or 6 hours after opening until closing time	✓ $N'(t) = 3t^2 - 24t + 36$ ✓ $N'(t) \geq 0$ ✓ $(t - 6)(t - 2)$ ✓ $t \geq 6$ ✓ $t \leq 2$ (5)
10.3	Minimum turning point at $t = 6$ hours after opening	✓ $t = 6$ (1) [7]

QUESTION 11

11.1.1	 OR $P(M \text{ or } N) = P(M) + P(N) - P(M \text{ and } N)$ $0,6 = 0,4 + 0,3 - P(M \text{ and } N)$ $P(M \text{ and } N) = 0,4 + 0,3 - 0,6$ $= 0,1$	✓ $P(N \text{ and } M') = 0,2$ ✓ $P(M \text{ and } N') = 0,3$ ✓ $P(N \text{ or } M') = 0,4$ ✓✓ $P(M \text{ and } N) = 0,1$ (5)
11.1.2	$LH = P(M \text{ and } N)$ $= 0,1$ $RH = P(M) \times P(N)$ $= (0,4)(0,3)$ $= 0,12$ $\therefore P(M \text{ and } N) \neq P(M) \times P(N)$ $\therefore M$ and N is not independent	✓ $P(M \text{ and } N) = 0,1$ ✓ $P(M) \cdot P(N) = (0,4)(0,3)$ ✓ $0,12$ ✓ $P(M \text{ and } N) \neq P(M) \times P(N)$ ✓ No, not independent (5)

11.2	$1 \times 8 \times 7 \times 6 \times 2$ $= 672$ OR $(1 \times 8 \times 7 \times 6 \times 1) + (1 \times 8 \times 7 \times 6 \times 1)$ $= 672$	$\checkmark 1 \times$ $\checkmark 8 \times 7 \times 6$ $\checkmark \times 2$ $\checkmark 672$ $\checkmark 1 \times$ $\checkmark 8 \times 7 \times 6 \times 1$ $\checkmark + (1 \times 8 \times 7 \times 6 \times 1)$ $\checkmark 672$	 (4) (4) [14]
		TOTAL:	150