




eduresourceza.com



GAUTENG PROVINCE
EDUCATION
REPUBLIC OF SOUTH AFRICA

**GAUTENG DEPARTMENT OF EDUCATION
PREPARATORY EXAMINATIONS
2015**

MEMORANDUM

SUBJECT:	PHYSICAL SCIENCES P2 (10842)
-----------------	-------------------------------------

GAUTENG DEPARTMENT OF EDUCATION
PREPARATORY EXAMINATIONS

PHYSICAL SCIENCES
(Second Paper)

MEMORANDUM

QUESTION 1


- | | | |
|------|-----|-----|
| 1.1 | C✓✓ | (2) |
| 1.2 | B✓✓ | (2) |
| 1.3 | D✓✓ | (2) |
| 1.4 | C✓✓ | (2) |
| 1.5 | B✓✓ | (2) |
| 1.6 | A✓✓ | (2) |
| 1.7 | A✓✓ | (2) |
| 1.8 | A✓✓ | (2) |
| 1.9 | B✓✓ | (2) |
| 1.10 | C✓✓ | (2) |

[20]

QUESTION 2

- 2.1.1 C ✓ (1)
- 2.1.2 A ✓ (1)
- 2.2.1 Butanal ✓ (1)
- 2.2.2 2-methylbutan-2-ol ✓ (2)
OR 2-methyl-2-butanol
- 2.3.1 Water / H₂O ✓ (1)
- 2.3.2

- Correct functional group ✓
- Methyl group on the second carbon ✓
- Whole structure ✓

Notes:

- Condensed or semi-structural formula: $\frac{1}{3}$
- Molecular formula: $\frac{0}{3}$

(3)


- 2.3.3 Addition / Hydration✓ (1)
- 2.4.1 Process by which molecules of two monomers with different functional groups✓ react with the elimination/loss of water molecules. ✓ (2)
- 2.4.2 F✓ (1)
- [13]

QUESTION 3

- 3.1.1 Organic compounds having the same molecular formula✓ with the same functional group situated at different positions. ✓ or same side chain at different positions
OR
Organic compounds with the same molecular formula, ✓ but different positions of the side chain / functional groups. ✓

(2)


3.1.2



(1)

But -1-yne ✓ /1-butyne ✓

(1)




(1)

But-2-yne ✓ /2-butyne

(1)

3.2.1



Reactants✓ Products✓✓

(3)

Condensed/semi-structural formulae:	Max: $\frac{2}{3}$
Molecular formula;	$\frac{0}{3}$
Any additional reactant or products:	Max: $\frac{2}{3}$

3.2.2 Propene✓ and methylpropane ✓

(2)

3.3.1 The temperature ✓ at which the vapour pressure of a liquid is equal to the external/ atmospheric pressure ✓ ORThe temperature ✓ at which a liquid changes phase into the gaseous phase. $\frac{1}{2}$

(2)

3.3.2 Propan -1-ol has only one hydrogen bond ✓ between its molecules.Thus, propan – 1 – ol has weaker intermolecular forces between its molecules. ✓ ∴ Less energy is needed to overcome these intermolecular forces, ✓ thus propan – 1 – ol has a lower boiling point. ORPropanoic acid has two hydrogen bonds ✓ between its molecules.The intermolecular forces between the propanoic acid molecules are thus stronger ✓ ∴ more energy is needed to overcome these forces, ✓

thus propanoic acid has a higher boiling point.

(3)


[16]

QUESTION 4

4.1.1 Q: Substitution✓/halogenation/chlorination (1)

4.1.2 R: Elimination/ dehydrohalogenation/ dehydrobromination✓ (1)

4.2.1



- Functional group
- Whole structure correct

Accept -OH
condensed in
structural formula

(2)


Condensed/semi – structural formulae	Max: 1/2
Molecular formula	0/2

4.2.2 Propan – 2 – ol ✓ (1)

2-propanol

4.3.1 Hydrogen bromide ✓ (1)

4.3.2



• Functional group✓ • Whole structure correct✓	
Accept HBr Ignore: ⇌	
• Condensed/semi-structural formulae	Max: 3/4
• Molecular formula	0/4
Any additional reactant or products: Max: 3/4	Everything correct, <i>wrong balancing</i> Max: 3/4

(4)
[10]

QUESTION 5

5.1.1 Surface area/state of division/reaction surface✓ (1)

5.1.2 There are more particles available for reaction.✓ /more collisions.
There are a greater number of effective collisions✓per unit of time✓
 ∴ reaction rate increases (3)

5.1.3 Mass of 15% of original 2 g=0,3 g✓
 Number of moles of CaCO₃ 0,3 g:

$$n = \frac{m}{M}$$

$$= \frac{0,3}{100}✓$$

$$= 0,003 \text{ mol.}$$

n_{CaCO₃}: n_{HCl}

1 : 2

0,003: 0,006✓

$$c_{\text{HCl}} = \frac{n}{V}$$

$$0,2 = \frac{0,006}{V}$$

$$V = 0,03 \text{ dm}^3 ✓$$

✓ for both equations

5.2.1 Concentration✓ (1)


5.2.2 Equal to ✓ (1)

5.2.3 The same number of moles/mass/ number of particles of CaCO₃ ✓ (produce same number of moles of carbon dioxide). CaCO₃ is the limiting agent/Acid is in excess. (1)

5.3.1 (Experiment) 6✓

(1)

5.3.2



Criteria for graph		Mark
Correct shape for both graphs		✓
Graph of experiment 5 reaches a higher maximum/peak value than experiment 6.		✓
Graph of experiment 6 shows more molecules of higher kinetic energy		✓

(3)

If no labels on axes	max:2/3
----------------------	---------

5.4.1 exothermic✓

(1)

5.4.2 The energy/potential energy of the products is lower than that of the reactants ✓

OR

(1)

More energy released than absorbed✓

OR

[18]

Energy is released

QUESTION 6

6.1 CALCULATIONS USING NUMBER OF MOLES:

Mark allocation:

- USING formula $n = \frac{3,01 \times 10^{23}}{6,02 \times 10^{23}}$ ✓
- Change in $n(\text{NO}) = \text{Initial mol} - \text{equilibrium mol}$ ✓
- Using ratio $\text{NO(g)} : \text{O}_2(\text{g}) : \text{NO}_2(\text{g}) = 2 : 1 : 2$ ✓ to determine change in $n(\text{O}_2)$ and change in $n(\text{NO}_2)$
- Equilibrium $[\text{NO}] = \text{equilibrium } n(\text{NO}) / 0,05$
 Equilibrium $[\text{O}_2] = \text{equilibrium } n(\text{O}_2) / 0,05$
 Equilibrium $[\text{NO}_2] = \text{equilibrium } n(\text{NO}_2) / 0,05$ } ✓
- Correct K_c expression (formulae in []) ✓
- Substitution of concentrations and K_c value into K_c expression ✓
- $[\text{O}_2]$ at equilibrium ✓
- $n(\text{O}_2)$ at equilibrium ($\times 0,05$) ✓
- $n(\text{O}_2)$ initial = n at equilibrium + n change ✓

OPTION 1

$$n(\text{NO}) = \frac{3,01 \times 10^{23}}{6,02 \times 10^{23}} = 5 \times 10^{-3} \text{ mol}$$

	NO(g)	O ₂ (g)	NO ₂ (g)
Initial moles	1	0,5006 ✓	0
Change in moles	0,995 ✓	0,4975	0,995
Equilibrium moles	0,005	0,0031 ✓	0,995
Equilibrium concentration	0,1	0,0615	19,9

✓ mole ratio

✓ divide by 0,05

$$K_c = \frac{[\text{NO}_2]^2}{[\text{NO}]^2 [\text{O}_2]}$$

$$6,44 \times 10^5 = \frac{(19,9)^2}{(0,1)^2 [\text{O}_2]}$$

$$[\text{O}_2] = 0,0615 \text{ mol} \cdot \text{dm}^{-3}$$

[9]

- 6.2 Large K_c means a high yield of products compared to reactants✓ (1)
- 6.3 Darker✓ (1)
- 6.4 Temperature is decreased and the reaction will adjust to resist the change. The exothermic✓ reaction is favoured thus the forward reaction✓ is favoured and the contents become darker.✓ (3)
- 6.5 No effect ✓ (1)
- [15]

QUESTION 7

- 7.1 Brønsted-Lowry base accepts a proton✓✓ (2)
- 7.2.1 $\text{pH} = -\log[\text{H}_3\text{O}^+]$ ✓
 $= -\log(0,12)$ ✓
 $= 0,92$ ✓ (3)
- 7.2.2 Cl^- ✓ (1)
- 7.3.1 It can act as an acid or a base/ it can accept or donate an H^+ ion✓ (1)
- 7.3.2 $\text{HSO}_4^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{SO}_4^{2-}$
 ✓ reactants ✓ products ✓balancing
- Ignore → (3)
- 7.4 Drops of water will change the concentration of the acid/ make it more dilute✓ (1)
- 7.5.1 Option 1:
- $$n = \frac{m}{M}$$
- $$= \frac{8,0}{56}$$
- $$= 0,14 \text{ mol}$$
- $$c = \frac{n}{V}$$
- $$= \frac{0,14}{0,25}$$
- $$= 0,57 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$
- Option 2:
- $$c = \frac{m}{MV}$$
- $$= \frac{8,0}{56 \times 0,25}$$
- $$= 0,57 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$
- (3)

7.5.2 $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$ ✓
(dilute acid)

$$\frac{c_a(40)}{(0,57)(25)} \checkmark = \frac{1}{2} \checkmark$$

OR

$$c_a = \frac{1 \times 0,57 \times 25 \checkmark}{2 \times 40} \checkmark$$

$$= 0,178 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

$$\text{OR} \quad c_a = \frac{1 \times 0,57 \times 0,025 \checkmark}{2 \times 0,04} \checkmark$$

$$= 0,178 \text{ mol} \cdot \text{dm}^{-3} \quad (4)$$

7.5.3 **Concentrated acid**

$$n_1 \text{ (moles concentrated)} = n_2 \text{ (moles dilute)}$$

✓

$$c_1 \times V_1 = c_2 \times V_2$$

$$c_1 = \frac{0,178 \times (490 + 10) \checkmark}{10} \checkmark$$

$$= 8,9 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

OR

$$c_1 = \frac{0,178 \times 0,5 \checkmark}{0,01} \checkmark$$

$$= 8,9 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

(4)
[22]

QUESTION 8

8.1 Electrical energy to chemical energy ✓ (1)

8.2 Cathode ✓

At cathode, Ag^+ / silver ions gain electrons ✓ and are, reduced to Ag/ silver metal ✓




(3)

8.3.1 Ag/silver ✓

(1)

8.3.2 $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}$ ✓✓

(2)



8.4.1 Electrode X becomes eroded/ smaller/ thinner. ✓ (1)

8.4.2 A (silver) layer forms on the medal. ✓ (1)

8.5 The rate at which Ag^+ / silver ions are reduced at the cathode is equal to rate at which Ag metal/ silver is oxidised at the anode. ✓✓

(2)
[11]

QUESTION 9

9.1 Galvanic✓ cell (1)

9.2 There will no reading/ The reading will be zero/ 0V✓ (1)

9.3 Temperature ✓ and the initial concentrations of the electrolytes ✓

But cannot give 25 °C and 1 mol·dm⁻³ because the investigation is not carried out under standard conditions because the emf values are not for standard conditions. (2)

9.4 The different type of metal/the different half-cells✓ (1)

9.5.1 Voltmeter's terminals✓ have been connected incorrectly✓
OR
Incorrect connection ✓ (+ to anode and – to cathode) ✓ (2)

9.5.2 Aluminium is a stronger reducing agent than zinc✓ and zinc is a stronger reducing agent than copper.✓ (2)

9.6.1 Aluminium/ Al✓ (1)

9.6.2 Zinc/ Zn ✓ (1)

9.7 $2\text{Al}(\text{s}) + 3\text{Zn}^{2+}(\text{aq}) \longrightarrow 2\text{Al}^{3+}(\text{aq}) + 3\text{Zn}(\text{s})$ ✓ balancing ✓

Reactants✓ Products✓ Balancing✓

Ignore ⇌

Marking rule 3.9:

If only a reactant(s) followed by an arrow, or only a product(s) preceded by an arrow, is/are written, marks may be awarded for the reactant(s) or product(s). If only a reactant(s) or only a product(s) are written, without an arrow, no marks are awarded for the reactant(s) or product(s).

Examples: $\text{N}_2 + 3\text{H}_2 \longrightarrow 2\text{NH}_3$ ✓ bal. ✓

$\text{N}_2 + \text{H}_2 \longrightarrow$ ✓ $\frac{1}{3}$

$\longrightarrow \text{NH}_3$ ✓ $\frac{1}{3}$

$\text{N}_2 + \text{H}_2$ $\frac{0}{3}$

NH_3 $\frac{0}{3}$

(3)
[14]

QUESTION 10

10.1.1 Iron oxide/iron/aluminium oxide✓ (1)

10.1.2 more collisions per unit time✓
greater surface area/greater number of exposed particles.✓ (2)

10.1.3 Increase the number of moles of N_2 and H_2 (reactants)✓ favours the forward reaction✓

OR

Increase pressure ✓ and thus favours the forward reaction✓ (2)

10.2.1 Calcium hydroxide increases the pH/alkalinity of the soil/ reduce the acidity of the soil.✓

Ammonium sulphate is a fertiliser. ✓ (2)

10.2.2 Ammonia gas is formed which will be released and **reduce** the concentration/number of moles/ of fertiliser.✓ (1)

10.3 $n(NO) = \frac{m}{M} = \frac{720}{24} = 30 \text{ mol}$ ✓
 $n(HNO_3) = n(NO) = 30 \text{ mol}$ ✓ (3)

$m(HNO_3) = nM = (30)(63) = 1890 \text{ g}$ ✓ [11]

TOTAL: 150

eduresourceza.com