



eduresourceza.com

CAPE WINELANDS EDUCATION DISTRICT

CAPS

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY (P2)
MEMORANDUM SEPTEMBER 2015

MARKS 150

TIME 3 hours

This question paper consists of 9 pages.

QUESTION 1 [START ON A NEW PAGE]

- 1.1 A ✓✓ (2)
- 1.2 D ✓✓ (2)
- 1.3 C ✓✓ (2)
- 1.4 C ✓✓ (2)
- 1.5 D ✓✓ (2)
- 1.6 D ✓✓ (2)
- 1.7 B ✓✓ (2)
- 1.8 D ✓✓ (2)
- 1.9 A ✓✓ (2)
- 1.10 D ✓✓ (2)

[20]

QUESTION 2 [START ON A NEW PAGE]

- 2.1 Organic compounds that consist of hydrogen and carbon only. ✓✓ (2)
- 2.1.1 B ✓ (1)
- 2.2.2 C ✓. (1)
- 2.2.3 E ✓ (1)
- 2.3 Organic molecules with the same molecular formula, but different structural formulae ✓✓ (2)


- 2.4 2-methylprop-1-ene ✓✓ or methylpropene
(Double bond correctly named ✓ side chain correctly named ✓) (2)


Marking criteria

IF/INDIEN:

2 methylprop1 ene / 2 metielpropeen 1 een (1 mark out of two)

- 2.5 C_4H_8 ✓ (1)

- 2.6  functional group ✓ rest of molecule ✓. (2)

- 2.7  ✓ if I H H no marks (1)

- 2.8 Butanoic acid ✓ / butanoesuur (1)

- 2.9 unsaturated ✓ Compounds with one or more multiple bonds between C atoms ✓✓
in their hydrocarbon chains. (3)

ANY ONE/ENIGE EEN:

- It does not ONLY have single bonds.
- It does not have single bonds between all C atoms.
- It has double bonds between C atoms.
- It does not contain the maximum number of H atoms bonded to C atoms.
- Each C atom in B is not bonded to four other atoms.

- 2.10 alkenes ✓ (1)

[18]

QUESTION 3 [START ON A NEW PAGE]

- 3.1.1 Different functional group/homologous series ✓
Verskillende funksionele groepe/homoloe reeks (1)
- 3.1.2 Boiling point/Kookpunt ✓ (1)
- 3.2 Higher ✓ Between the particles of propan-1-ol there are hydrogen bonds and between the particles of propanal there are London forces (induced dipole forces) (momentele dipool kragte) ✓. Hydrogen bonds are stronger than London forces ✓
Hydrogen bonds require more energy to overcome than London forces. ✓ (4)
- 3.3.1 A propanal ✓ (1)
- 3.3.2 B - propan-1-ol ✓ (1)
- 3.3.3 C ethanoic acid ✓ (1)
- 3.4 HIGHER, ✓ the van der Waals forces increases with increasing molecular mass ✓
or the longer the carbon chain/greater the surface area, the greater stronger the London forces will become. (2)

[11]

QUESTION 4 [START ON A NEW PAGE]

- 4.1.1
- $$\begin{array}{c} \text{H} \\ | \\ \text{C} = \text{C} - \text{C} - \text{H} \\ | \quad | \quad | \\ \text{H} \quad \text{H} \quad \text{H} \end{array} + \text{H} - \text{O} - \text{H} \longrightarrow \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ | \quad | \quad | \\ \text{H} - \text{C} - \text{C} - \text{C} - \text{H} \\ | \quad | \quad | \\ \text{H} \quad \text{O} \quad \text{H} \end{array}$$
- (4)
- 4.1.2 propan-2-ol/2-propanol (1)
- 4.1.3 a. substitution / hydrolysis ✓ (1)
b. heat ✓
dilute NaOH or KOH (under reflux) ✓ (2)
- 4.2.1 esterification ✓ (1)
- 4.2.2 hydrogen sulfate / H₂SO₄ ✓ (1)

4.2.3 $M[C_3H_6O_2] = 74$
 $n = \frac{m}{M_r} \checkmark = \frac{68,88}{74} \checkmark = 0,93 \text{ mol}$
 0,93 mol ester was delivered by 0,93 mol methanol ✓
 $M[CH_4O] = 32$
 $0,93 \times 32 \checkmark = 29,78 \text{ g}$
 % purity: $\frac{29,79}{50} \times 100 = 59,57\% \checkmark$

- One mark for formula
- One mark for correct substitution of M
- One mark for the USE of the molar relationship
- One mark for calculating the mass of methanol formed.
- One mark for calculating percentage purity.

(5)
[15]

QUESTION 5 [START ON A NEW PAGE]

- 5.1 The zinc becomes smaller ✓
 or gas bubbles forms ✓
 or the test tube heats up
 (any two) (2)
- 5.2 The change in concentration of reactants or products per unit time. ✓ ✓
 also except change in mol/volume/mass of reactants /products per unit time (2)
- 5.3 $\text{rate} = - \frac{\Delta c}{\Delta t} \checkmark = - \left(\frac{0,77 - 2 \checkmark}{45 - 0 \checkmark} \right) = 0,027 \text{ mol.dm}^{-3} \cdot \text{s}^{-1} \checkmark$ (4)
- 5.4
- 5.4.1 increase ✓ (1)
- 5.4.2 stay the same ✓ (1)
- 5.4.3 increase ✓ (1)

[11]

QUESTION 6 [START ON A NEW PAGE]

- 6.1 It is an equilibrium where the rate of the forward reaction equals the rate of the reverse reaction ✓✓ (2)
- 6.2 Between t_1 and t_2 ✓ and between t_3 and t_4 ✓ (2)
- 6.3 reverse ✓ (1)
- 6.4 EXOTHERMIC ✓ (1)
- 6.5 - $[CO]$ and $[O_2]$ decrease and $[CO_2]$ increases ✓
 - This means the forward reaction is favoured. ✓
 - decrease in temperature, favours exothermic reaction, therefore reaction is exothermic. ✓ (3)
- 6.6 The concentration of CO_2 was suddenly increased (by adding more CO_2 gas) ✓
 OR $[CO_2]$ increases / CO_2 is added. (1)
- 6.7 Remains the same ✓ (1)

6.8 Option 1

$$n(CO) \text{ initially: } n = \frac{m}{M} = \frac{63}{28} = 2,25 \text{ mol}$$

$$n(O_2) \text{ initially } n = \frac{m}{M} = \frac{9,11}{32} = 0,28 \text{ mol}$$

one mark for both values ✓

At equilibrium:

$[CO_2]$ is $0,15 \text{ mol} \cdot \text{dm}^{-3}$ (given), therefore

$$n(CO_2) \text{ formed: } n = cV = 0,15 \times 2 = 0,3 \text{ mol } ✓$$

$$n(O_2) \text{ reacted: } 0,15 ✓$$

$$n(CO) \text{ reacted: } 0,3 ✓$$

$$n(CO) \text{ at equilibrium: } 2,25 - 0,3 = 1,95 ✓$$

$$n(O_2) \text{ at equilibrium: } 0,28 - 0,15 = 0,13 ✓$$

Calculate concentration: ✓

$$K_c = \frac{[CO_2]^2}{[CO]^2[O_2]} ✓$$

$$= \frac{(0,15)^2}{(0,975)^2(0,065)} ✓$$

$$= 0,36 ✓$$

OPTION 2:

	2CO	O₂	2CO₂
Initial	2,25	0,28	0 (✓ one mark for all values in row)
Reacted/ formed	0,3✓	0,15✓	0,3
At equilibrium	1,95✓	0,13✓	0,3✓
Equilibrium concentration	0,975	0,065	0,15 (✓ one mark for all values in row)

$$K_c = \frac{[CO_2]^2}{[CO]^2[O_2]} \checkmark$$

$$= \frac{(0,15)^2}{(0,975)^2(0,065)} \checkmark$$

$$= 0,36 \checkmark$$

(10)

[21]

QUESTION 7

7.1.1 The point where the indicator changes colour.. ✓ ✓ (2)

7.1.2 An acid that is able to donate two protons (H⁺) per unit ✓ (1)

7.1.3 H₂SO₄ ionises completely in water to form a high concentration of H₃O⁺ ions. ✓ ✓ (2)

7.1.4 20 cm³ ✓ (1)

7.1.5 $n = cV$ ✓
 $= 0,2 \times 0,05$ ✓
 $= 0,01$ mole
 0,01 mol NaOH reacts with 0,005 mole H₂SO₄ ✓

$$c = \frac{n}{v}$$

$$= \frac{0,005}{0,02} \checkmark$$

$$= 0,25 \text{ mol} \cdot \text{dm}^{-3} \checkmark$$

(5)

- 7.2.1 It can act as either an acid or a base. ✓ ✓ (2)
- 7.2.2 $10^{-7} \text{ mol} \cdot \text{dm}^{-3}$ ✓ (1)
- 7.2.3 $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$ ✓ ✓ (2)
- 7.3.1 Decrease ✓ (1)
- 7.3.2 According to Le Chatelier, an increase in temperature will favour the forward reaction. ✓, . That means the $[\text{H}_3\text{O}^+]$ increases and therefore the pH will decrease. ✓ (2)

[19]


QUESTION 8 [START ON A NEW PAGE]

- 8.1.1 chemical (potential) energy ✓ to electrical energy ✓ (2)
- 8.1.2 a solution/dissolved substance that conducts electricity through the movement of ions ✓ ✓ (2)
- 8.1.3 B ✓ (1)
- 8.1.4 $\text{Ni}^{2+} + 2\text{e}^- \rightarrow \text{Ni}$ ✓ ✓ (if double arrow only one mark)
- Notes**
- | | | | |
|---|----------------------------|---|----------------------------|
| $\text{Ni} \leftarrow \text{Ni}^{2+} + 2\text{e}^-$ | $\left(\frac{2}{2}\right)$ | $\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$ | $\left(\frac{1}{2}\right)$ |
| $\text{Ni} \rightleftharpoons \text{Ni}^{2+} + 2\text{e}^-$ | $\left(\frac{0}{2}\right)$ | $\text{Ni}^{2+} + 2\text{e}^- \leftarrow \text{Ni}$ | $\left(\frac{0}{2}\right)$ |
- (2)
- 8.1.5 $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$ ✓
- $= -0,27 \text{ V} - (-2,36 \text{ V})$ ✓
- $= 2,09 \text{ V}$ ✓

Notes

Accept any other correct formula from the data sheet
Any other formula using unconventional abbreviations, e.g. $E^\circ_{\text{cell}} = E^\circ_{\text{OA}} - E^\circ_{\text{RA}}$
followed by correct substitutions: $\left(\frac{3}{4}\right)$

Option 2



(4)

CAPS

- 8.1.6 From B to A ✓ (1)
- 8.1.7 Anions moves to Ni-half cell (A) ✓
Cations moves to mg-half cell (B) ✓
Ensures electrical neutrality to ensure flow of electrons in the external circuit ✓ (3)
- 8.1.8 $\text{Mg(s)}|\text{Mg}^{2+}(\text{aq})||\text{Ni}^{2+}(\text{aq})|\text{Ni(s)}$ oxidation half cell correct ✓
reduction half cell correct ✓
saltbridge ✓ (3)
If phases not included, do not subtract marks.
- 8.2.1 $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}$ ✓ ✓ (if double arrow only one mark)
Notes
 $\text{Al} \leftarrow \text{Al}^{3+} + 3\text{e}^- \quad \left(\frac{2}{2}\right) \quad \text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al} \quad \left(\frac{1}{2}\right)$
 $\text{Al} \rightleftharpoons \text{Al}^{3+} + 3\text{e}^- \quad \left(\frac{0}{2}\right) \quad \text{Al}^{3+} + 3\text{e}^- \leftarrow \text{Al} \quad \left(\frac{0}{2}\right)$ (2)
- 8.2.2 Electrode B (carbon) ✓ (1)
- 8.2.3 Al^{3+} ✓ or/of Al_2O_3 (1)

[22]**QUESTION 9 [START ON A NEW PAGE]**

- 9.1 Nitrogen/stikstof (N_2) ✓ (Not N) (1)
- 9.2 $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ ✓ balancing ✓ (Ignore arrow) (3)
- 9.3 Ammonia/ammoniak ✓ (1)
- 9.4 $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g})$ ✓ balancing ✓ (3)
- 9.5 ammonium nitrate/ammoniumnitraat ✓ (1)
- 9.6 Eutrophication/eutrifikasie (1)
- 9.7 $40/100$ (or $0,4$) $\times 1,5$ ✓ = $0,6\text{ kg}$ ✓ fertilizer in bag.
 $1/1 \times 0,6$ ✓ = $0,6\text{ kg}$ ✓ (600 g) Nitrogen is present in fertilizer
(OR only nitrogen/no other fertilizer present, $0,6\text{ kg}$ nitrogen) (4)

[14]**TOTAL 150**