



eduresourceza.com



NATIONAL SENIOR CERTIFICATE EXAMINATION
NOVEMBER 2018

EQUINE STUDIES
MARKING GUIDELINES

Time: 3 hours

200 marks

These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.

The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.

SECTION A

QUESTION 1

- 1.1 Give term described below
 - 1.1.1 Fallopian tube OR oviduct
 - 1.1.2 Prebiotic
 - 1.1.3 Molar
 - 1.1.4 Prostaglandin
 - 1.1.5 Lymphangitis
- 1.2 Give the definition of the following
 - 1.2.1 Small living particle which infects and causes disease, not visible with ordinary microscope.
 - 1.2.2 Bleeding from nose
 - 1.2.3 Sugar substance in food, simple and complex
 - 1.2.4 Building block of protein.
 - 1.2.5 Connect artery and vein, very small.
- 1.3
 - 1.3.1 False. Thrush produces tarry black discharge from the frog Sulci
 - 1.3.2 True
 - 1.3.3 True
 - 1.3.4 False. Navicular is a disease of the distal sesamoid bone
 - 1.3.5 False. Sheared heels is when the bulbs of each heel move independently of each other OR contracted heels.
 - 1.3.6 False. Side bone is seen in older, heavy, draught-type horses.

QUESTION 2

- 2.1 2.1.1 1. C
 2. G
 3. F
 4. B
 5. E
 6. H
 7. A
- 2.1.2 Tying up, muscle sprain/tear, muscle fatigue, atrophy. (any two)
- 2.2 2.2.1 Suspensory ligament, DDFT, SDFT (correct order)
- 2.2.2 Rest to reduce swelling to start healing process prevent any further damage. Walk to start aligning tendon fibres so heal straight and stronger. Increase blood flow for healing.
- 2.3 1. capped elbow
 2. capped knee / carpalis
 3. wind galls/wind puffs
 4. bog spavin
 5. capped hock
 6. thoroughpin

QUESTION 3

- 3.1 3.1.1 Alveoli of the lung showing gaseous exchange
- 3.1.2 Red blood cell
- 3.1.3 Haemoglobin
- 3.1.4 Carry oxygen
- 3.1.5 Oxygen
- 3.1.6 Carbon dioxide
- 3.1.7 High concentration to a low concentration
- 3.1.8 Increased respiration to move more air, increased heart rate to increase blood flow and therefore amount of oxygen to the muscles, spleen contracts to release more RBC so more oxygen can be transported
- 3.1.9 Alveolus wall
- 3.1.10 Capillary wall
- 3.1.11 They are only one cell thick and therefore make it easier for gases to diffuse across a smaller distance.

SECTION B

QUESTION 4

4.1 Looking at Figure 3 on endocrine control of reproduction

4.1.1 Sertoli – testicle interstitial tissue; Leydig – testical seminiferous tubules

4.1.2 Libido, crest formation, deep voice, aggressive behaviour. Spermatogenesis, increase in testicular size

4.1.3 Follicle-stimulating hormone

4.1.4 Luteinising hormone luteinises the corpus luteum. Triggers ovulation

4.2 Looking at Figure 4

4.2.1 Seasons are different / opposite to ours in southern hemisphere

4.2.2 Testosterone is needed to stimulate sperm production.

4.2.3 April till August as sperm count is highest and mares in season as it is summer time in northern hemisphere so highest chance of getting mare pregnant.

QUESTION 5

- 5.1 12% protein content
- 5.2 No one feed fits all, thriving and poorly underweight horses will require higher-fat meal overweight horses will require restricted-energy feed with higher roughage, horse may be lazy and need more energy as this food is good for a hot horse, horse may be in hard work and require higher protein, age of horse will affect protein and energy requirements young horse requires more protein to grow than adult fully grown horse older horses have specific requirements want to feed as little as possible may need to feed large amounts to meet energy requirements.
- 5.3 $120 \text{ g/kg} \times 4 = 480 \text{ g protein}$
- 5.4 5.4.1 Add lucerne as high in Ca or supplement with any Ca-containing feed.
5.4.2 May get miller's disease.
- 5.5 Raw maize not easily digested, undigested starch spills into hind gut, digested there into lactic acid causes horse to be hot, no longer "cool and perform".
- 5.6 Micronised is infrared heated, therefore starch more easily digested and available to horse, will not look like whole kernel, often flaked or in pellets as tiny pieces.
- 5.7 Keeps horse calm and keeps weight on, slow-release energy for horse, higher energy density therefore feed less.
- 5.8 Scavenging free radicals increase in immunity, prevents oxidation which causes damage which causes ageing and inflammation
- 5.9 Salt forms part of electrolytes which are important for chemical reactions and specific functions in horses; horses lose salt when sweating during work, therefore need to replace. Nerve impulses, increases need to drink water
- 5.10 Decrease concentrate increase roughage
- 5.11 Still needs to maintain body and extra to heal body – reason for box rest.
- 5.12 Only change in third trimester as this is when biggest growth of the foetus occurs therefore higher protein or increased feed amount will be required, give smaller feeds more often as foetus takes up room in abdomen putting pressure on stomach, good-quality hay and grazing free access to fresh, clean water
- 5.13 Move from bully horse, give hay first, feed on schedule, not late, large, smooth rocks in feed, smaller amounts more often

QUESTION 6

- 6.1 Viral infection that affects blood vessels and lungs
- 6.2 Infected, protection, surveillance and free
- 6.3 No, only vaccinated in infected and protection zone, horses in surveillance zone can request permission to vaccinate, free zone. No vaccinations but may request permission to vaccinate June October
- 6.4 Need to stay AHS free so export of horses possible, very important for racing industry, financial implications
- 6.5 June to October, two injections, three weeks apart
- 6.6 Decrease chance of vaccine mutation and spread by midges as live attenuated virus in vaccine can infect unvaccinated horses through midges
- 6.7 Vaccinations before movement, in a passport, health check by vet, movement permit by local state vet, inform/prenotification Boland state vet, possible quarantine stop over
- 6.8 Take temperature twice daily two days pre-trip and five days post-trip to check for travel sickness change feed week before trip to increase water in gut possibly with balanced bran, feed and water on trip, ad-lib hay as low as possible, stop every 5 hours to rest and water horse, depend on weather if blanket or sweat sheet used, poll guard and leg bandages more supportive than boxing boots for long distance, overreach boots, tail guard. (pre-travel, feed, water, stops, dress)
- 6.9 Right side due to camber of the road
- 6.10 Fever, anorexia, cough respiratory distress weight loss
- 6.11 Elbows turned out, head and neck extended and down, heaving or abdominal component, possible wheezing audible respiration (any three)
- 6.12 Fluid in alveolar where exchange of oxygen takes place is filled with fluid, inhibits oxygen and CO₂ exchange leads to decreased O₂ in blood therefore increased respiratory rate to exchange where no fluid is

SECTION C

QUESTION 7

- 7.1 Height accurate, width accurate, oval, small "mouth"
- 7.2 Anterior towards the head, posterior towards the tail
- 7.3 The eggs shed by adult flukes are expelled with the faeces. Once outside the host, the larvae called **miracidia** hatch out of the eggs in a few days. These larvae can survive for weeks without a host provided there is enough humidity. They die quickly in a dry environment. Miracidia can swim and penetrate actively into the snails where they develop successively to **sporocysts**, **rediae** and **cercariae**, the usual larval stages of most fluke species.
Mature cercariae leave the snail, attach to the vegetation and produce **cysts**, the so-called **metacercariae**, which are infective for the final host. Horses become infected when grazing on contaminated pastures. Once a final host ingests metacercariae while grazing, they migrate to their predilection sites.
- 7.4 Anaemia
- 7.5 Eggs seen in stool under microscope or adults seen in gut on post-mortem
- 7.6 Zoonosis
- 7.7 Wrong diagnosis, incorrect drug, incorrect amount of drug, re-infestation (any three)
- 7.8 Remove from paddock with dam, fence dam, cut grass around dam, eradicate snails with chemicals or biocontrol, drain dam/water.
- 7.9 Strategic deworming after faecal float only deworm horses with high egg count.

Total: 200 marks